

TEMPLATE FOR RELEVANT LABEL PARTICULARS (RLPs)
(Veterinary Products)

Select appropriate:

- New Product (include all applicable RLPs) OR
 Variation (highlight instructions that are being varied). Approval no. of label being varied:
67576/56378

Signal heading:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING FOR ANIMAL TREATMENT ONLY
Product name:	PROGUARD™ Liquid Sheep Blowfly Treatment
Active constituent/s:	ACTIVE CONSTITUENT: 500 g/L CYROMAZINE
Statement of claims:	Protects long wool sheep for up to 14 weeks from strike by blowfly (<i>Lucilia cuprina</i>) including organophosphate resistant strains, when applied by jetting or dipping.
Net contents:	5 [7,10] LITRES
Directions for Use Heading:	READ DIRECTIONS FOR USE BEFORE USING THIS PRODUCT DIRECTIONS FOR USE
Restrains:	Restrains: DO NOT USE on sheep which are producing or may in the future produce milk that may be used or processed for human consumption. DO NOT USE on sheep with less than 6 weeks wool. DO NOT collect and re-use fluid which runs off.
Contraindications:	Contraindications: Should not be used on open wounds e.g. those resulting from marking or mulesing operations.
Precautions:	
Side effects:	



Dosage & administration:

Dosage and administration

Mixing

Shake well before use.

Settling may occur after storage for some weeks. Stir product or invert container several times before opening. Pour the liquid into the vat through a strainer to remove any dry particles or flakes which can occasionally occur under hot storage conditions. Stir well. This product is readily dispersible in both soft and hard water.

Dilution Rate

Dilution rate: 2mL per 1 Litre water

Examples: 400mL per 200 Litres
2L per 1000 Litres

Maintenance of Dips

1. Clean out dip, sump and spray lines thoroughly before charging. Disinfect with a registered bacteriostat at the manufacturer's recommended rate.
2. DO NOT USE old fouled wash. Shower dips should be emptied and cleaned out at the end of each day's dipping. Plunge dips should be emptied and cleaned out within 5 days of charging and more frequently if conditions favour post-dipping lameness infection, except in WA where plunge dips and sumps should be emptied daily.
3. Before use, stir well.

ANIMAL/PEST	TREATMENT	INITIAL CHARGE	TOPPING UP	CRITICAL COMMENTS
Sheep Blowfly Control (<i>Lucilia cuprina</i>)	Jetting	2L per 1000L	-	<ol style="list-style-type: none">1. Jet sheep at onset of fly wave.2. Apply 2 to 4L of jetting fluid per sheep, depending on sheep size and wool length.3. Ensure saturation to the skin.4. Pay particular attention to the treatment of the crutch, pizzles and polls.5. If used in jetting races a reduced period of protection may result.6. Heavy rain following the application of this product could diminish the period of protection.

		Dipping (Plunge, Shower and Constant Replenishment)	2L per 1000L	2L per 1000L	<ol style="list-style-type: none"> 1. Dip sheep at onset of fly wave. 2. Thoroughly saturate sheep. When using a shower dip, run top and bottom sprays independently so as to ensure maximum pressure and therefore penetration. 3. This is a non-stripping dip so there is no need to reinforce during dipping. 4. Dipping sheep heavily infested with grass seed is not recommended. 5. Heavy rain following the application of this product could diminish the period of protection.
		Dressing	2L per 1000L	-	<ol style="list-style-type: none"> 1. Clip the wool surrounding the strike, treat the struck area and the surrounding wool with a recommended knock down product then thoroughly jet the sheep with ProGuard. 2. This product is a slow killing larvicide and when used on its own, larvae may take 3 to 4 days to die.
General directions:	<p>General Directions</p> <p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p> <p>This product is designed for use on sheep with at least 6 weeks wool growth (long wool).</p>				

	<p>Jetting Jet the sheep at onset of fly wave with efficient equipment which is able to maintain the pressure of about 550 kPa. Use 300 to 400 kPa for lambs and ewes in lamb.</p> <p>Bacteriostat This product does not contain a bacteriostat. For control of post-dipping infections a registered bacteriostat should be added at the manufacturer's recommended rate.</p>
Withholding Period/s:	<p>WITHHOLDING PERIODS: MEAT: DO NOT USE less than 7 days before slaughter for human consumption. MILK: DO NOT USE on sheep which are producing or may in the future produce milk which may be used or processed for human consumption. WOOL HARVEST INTERVAL: DO NOT USE less than 2 months before shearing or fibre collection.</p>
Trade Advice:	<p>TRADE ADVICE:</p> <p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Landmark on 1300 255 347 or the APVMA website (www.apvma.gov.au/residues).</p> <p>Use of this product may result in wool residues that may not comply with European Union environmental quality standards.</p> <p>Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.</p>
Safety Directions:	<p>SAFETY DIRECTIONS</p> <p>May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the dip or jetting fluid, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. When using the prepared dip or jetting fluid, wear protective waterproof clothing, PVC or rubber apron, elbow-length PVC gloves, face shield and water resistant footwear. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.</p>
First Aid:	<p>FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.</p>
Additional user safety:	<p>ADDITIONAL USER SAFETY INFORMATION:</p> <p>Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling.</p> <p>Additional information is listed in the Material Safety Data Sheet.</p>
Environmental statements:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT DO NOT contaminate dams, ponds, waterways or drains with the chemical or used containers.</p>
Disposal:	<p>DISPOSAL This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered</p>

	place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.
Storage:	<p>Storage</p> <p>Keep out of reach of children. Store below 30°C (Room Temperature), in the original containers, tightly closed. Do not store for prolonged periods in direct sunlight.</p>
Name & address:	<p>Landmark Operations Limited</p> <p>24-26 Hydrive Close</p> <p>DANDENONG VIC 3175</p> <p>Customer Service 1300 255 347</p>

The following is for APVMA use only:

APVMA approval no.	APVMA No: 67576/0614
--------------------	-----------------------------