Product Name:GROCHEM DUSTING SULPHUR FUNGICIDE/MITICIDEAPVMA Approval No:67918/127503



Label Name:	GROCHEM DUSTING SULPHUR FUNGICIDE/MITICIDE
Signal Headings:	

Constituent Statements:	ACTIVE CONSTITUENT: 930 g/kg SULPHUR (S) - PRESENT AS POWDERED SULPHUR
Otatements.	

Mode of Action:		
	GROUP M2	FUNGICIDE

Statement of Claims:	For the control of powdery mildew of pumpkins, marrows, grapes and paw paws and for the control of citrus rust mite in citrus as per the Directions for Use Table.
----------------------	--

Net Contents: 25 kg

Restraints:	DO NOT apply to wet foliage.
Restraints:	DO NOT apply to wet tollage.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	NOT REQUIRED WHEN USED AS DIRECTED.

Trade Advice:	

	Apply the dust lightly and evenly to cover the foliage stems of the plant thoroughly. Rain will remove the dust deposit easily so that a fresh application is required after rain has fallen, as soon as the plants are dry. Crop Safety At temperatures over 24°C there is the possibility of leaf scorching if sensitive plants are treated, therefore reduced dosages are recommended during the hot dry months. Avoid dusting plants under windy conditions or when temperatures exceed 30°C.
--	---

Resistance Warning:	FUNGICIDE RESISTANCE WARNING
	GROUP M2 FUNGICIDE
	Grochem Dusting Sulphur Fungicide/Miticide is a member of the multi-site activity group of fungicides. For fungicide resistance management Grochem Dusting Sulphur Fungicide/ Miticide is a Group M2 fungicide. Some naturally occurring individual fungi resistant to the product and other Group M2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group M2 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Grochem Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

Precautions:	Caution Dust when the air is still. Avoid excessive clouds of the dust as sulphur is very irritating to the eyes.
--------------	---

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or watercourses with chemical or used containers. PROTECTION OF LIVESTOCK Low hazard to bees. May be applied to plants at any time.

Storage and Disposal:	KEEP OUT OF REACH OF CHILDREN Store in a closed original container in a dry, cool, well-ventilated area, out of direct sunlight. Shake and empty contents into spray tank. Do not dispose of undiluted chemicals on site. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product
	product.

Safety Directions:	
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: Australia 13 11 26; New Zealand 0800 764 766.

First Aid Warnings:

DIRECTIONS FOR USE

CROP	PEST	STATE	CRITICAL COMMENTS
Citrus	Citrus Rust Mite	Qld, NSW, Vic, SA & WA only	Apply from budburst onwards every 7 - 14 days.
Grapes	Powdery mildew	Qld, NSW, Vic, Tas, SA & WA only	
Paw Paws		Qld only	
Pumpkins, marrows		Qld, NSW, Vic, Tas, SA & WA only	Apply from seedling stage onwards every 7 - 14 days.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.