

Product Name: ACCENSI IPRODIONE 365 SE FUNGICIDE
APVMA Approval No: 68088/108372



Label Name:	ACCENSI IPRODIONE 365 SE FUNGICIDE
Signal Headings:	KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	365 g/L IPRODIONE
Mode of Action:	GROUP 2 FUNGICIDE
Statement of Claims:	A suspo-emulsion formulation with improved rainfastness (over other water based formulations) for the control of fungal diseases in recreational turf as specified in the directions for use.
Net Contents:	5-20L
Directions for Use:	
General Instructions:	<p>NOTE Accensi Iprodione 365 SE Fungicide may be very unstable in conditions where the pH is 7 or higher. It is therefore essential to check the pH of the spray solution before adding Accensi Iprodione 365 SE Fungicide. A suitable registered buffering agent may have to be added to bring pH down below 7.</p> <p>APPLICATION Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage.</p> <p>TURF NUTRITION This product acts very rapidly and where the turf is of a low nutritional status, a slight yellowing may be seen. This will be corrected by a nitrogenous fertiliser application. Where</p>

	<p>possible, it is recommended that the fertiliser program should precede the fungicide application.</p> <p>COMPATIBILITY The use of a surfactant or spray oil is not recommended with Accensi Iprodione 365 SE Fungicide as it may result in damage to sensitive plants. DO NOT mix with fertilisers. Mixtures with some fertilisers, e.g. urea, may cause foliar damage. As formulations of other manufacturers' products are beyond the control of Accensi Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.</p>
Resistance Warning:	<p>FUNGICIDE RESISTANCE STRATEGY</p> <p>GROUP 2 FUNGICIDE</p> <p>For fungicide resistance management Accensi Iprodione 365 SE Fungicide is a Group 2 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by the product and other Group 2 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Accensi Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant fungi.</p>
Precautions:	<p>DO NOT GRAZE TREATED AREAS OR FEED TURF CLIPPINGS FROM TREATED AREAS TO ANIMALS INCLUDING POULTRY</p>
Protections:	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p>
Storage and Disposal:	<p>Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product must not be burnt.</p>
Safety Directions:	<p>May irritate the eyes. Avoid contact with the eyes. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and water-proof gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.</p>
First Aid Instructions:	<p>If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.</p>

First Aid Warnings:	
---------------------	--

Directions for Use			
SITUATION	PEST	RATE	CRITICAL COMMENTS
Recreational Turf	Brown Patch (<i>Rhizoctonia solani</i>)	80 to 125 mL/4 to 150 L water/100 m ²	<p>Curative Program: Repeat treatment after 14 days. Use the higher rate under conditions of severe disease pressure.</p> <p>For preventative treatment: Apply as a monthly spray program. Repeat application at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. DO NOT apply more than 2 consecutive sprays of Accensi Iprodione 365 SE Fungicide or related dicarboximide fungicides unless tank mixed with a fungicide from a different activity group.</p>
	Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	80 mL/4 to 110 L of water/100 m ²	<p>For curative treatment: Repeat treatment in 14 days.</p>
	Winter Fusarium or Fusarium patch (<i>Fusarium rivale</i>) (NOT Qld)	125 mL/4 to 150 L of water/100 m ²	<p>For preventative treatment: Apply monthly. Repeat application at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. Do not apply more than two consecutive sprays of Accensi Iprodione 365 SE Fungicide or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.</p>
	Helminthosporium disease affecting leaf, crown and roots: (Black and white <i>Helminthosporium</i> spot <i>Bipolaris</i> spp. <i>Drechslera</i> spp. <i>Exserohilum</i> spp.)	65 mL/4 to 150 L of water/100 m ²	<p>For curative treatment: Apply as a foliar spray and repeat application after 7 to 14 days. For white spot symptoms on couch grass (<i>Cynodon dactylon</i>) only use higher water volumes of 100 to 150 L of water per 100m².</p> <p>For preventative treatment: Apply as a high volume foliar spray. Repeat application at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. Do not apply more than two consecutive sprays of Accensi Iprodione 365 SE Fungicide or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.