

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

APPARENT PILFERER 500 HERBICIDE

ACTIVE CONSTITUENT: 500 g/L PROPYZAMIDE

SOLVENT: 42 g/L ETHYLENE GLYCOL



GROUP	D	HERBICIDE
--------------	----------	------------------

For selective control of certain grasses and broad leaf weeds in canola, grain legumes, legume pasture, lettuce and turf according to the Directions for use Table

APVMA Approval Number 68267/122248

AIRR APPARENT Pty Ltd
15/16 Princes Street, Newport NSW 2106
M 0411 227 338
E enquiries@apparentag.com.au
www.apparentag.com.au

Contents: 5L, 10L, 20L, 110L, 1000L

Directions for use

Restraint: DO NOT apply more than once per season.

DO NOT apply to water logged soils.

DO NOT apply to land where runoff to aquatic areas or sensitive crops like cereals is likely.

CROP	WEEDS CONTROLLED	STATE	RATE L/HA	CRITICAL COMMENTS
Canola	Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue, Rats Tail Fescue	ALL STATES	1.0	Apply to moist soil prior to seeding only. Incorporate by sowing (IBS) with knife or blade points. A rainfall event (25mm) between application and sowing, or soon after sowing is required for best results. Avoid seeding practices that move treated soil back into the furrow. Heavy rainfall events that wash product back into the furrow can delay emergence and cause crop injury. DO NOT use on white sandy soils, or paddocks that have been recently mouldboard ploughed or spaded.
Grain Legumes (including lupins, chickpeas, lentils, field peas, faba beans, navy beans)				
Legume Pastures (including lucerne, clover, medics etc. grown for forage, hay or seed production)		NSW,ACT, TAS,SA, WA only	1.0-1.5	Apply when weeds are at post or pre-emergent stages. SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July.
	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, Seedling Sorrel, Blackberry Nightshade, Patterson's Curse, Seedling Wireweed		1.5-2.0	Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended

Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Ryegrass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	ALL STATES	4.5 in 330-550L of water	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
Lettuce (transplanted)				Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
		VIC only	2.3 in 300-550L of water	Apply on light sandy soils, low in organic matter.	
Turf and Lawns Common Couch	Winter grass	ALL STATES	1.2 (12mL/100m ²)	Rate/Bowling Green 37 x 37m 170mL	Apply during the autumn and winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Qld Blue Couch after 4 weeks. DO NOT RE-SEED FOR 60 DAYS AFTER TREATMENT. Use ONLY on established turf. DO NOT use on bluegrass, ryegrass, fescue or bent grass lawns.
Queensland Blue Couch		QLD only	0.6 (6 mL/100m ²)	85mL	
Buffalo Grass		WA only	1.2 (12 mL/100m ²)	170mL	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

CANOLA:

HARVEST – NOT REQUIRED WHEN USED AS DIRECTED

GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION

GRAIN LEGUMES:

HARVEST – NOT REQUIRED WHEN USED AS DIRECTED

GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION

ALL OTHER CROPS:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

MIXING

Prior to pouring, shake container vigorously, then add the required quantity of Apparent Pilferer 500 Herbicide to water in the spray vat while stirring or with agitators in motion.

DO NOT mix with other pesticides.

RESISTANT WEEDS WARNING

GROUP	D	HERBICIDE
--------------	----------	------------------

Apparent Pilferer 500 Herbicide is a member of the benzamide group of herbicides with inhibitors of microtubule assembly mode of action. For weed resistance management Apparent Pilferer 500 Herbicide is a Group D herbicide. Some naturally occurring weed biotypes resistant to Apparent Pilferer 500 Herbicide and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Pilferer 500 Herbicide or other Group D herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, AIRR Apparent Pty Ltd accepts no liability for any losses that may result from failure of Apparent Pilferer 500 Herbicide to control resistant weeds. Advice as to strategies and alternative treatments that can be used can be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or AIRR Apparent Pty Ltd representative.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

DO NOT graze sheep on treated areas or feed to poultry or pigs.

PROTECTION OF CROPS, NATIVE AND OTHER NONTARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping land or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Protect from frost.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in disposal pit specifically marked and set up for this purpose

clear of waterways, desirable vegetation and tree roots, in accordance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766