

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Burst 500 WG Herbicide

ACTIVE CONSTITUENT: 500 g/kg PROPYZAMIDE

GROUP	D	HERBICIDE
-------	---	-----------



For selective control of certain grasses and broad leaf weeds in Lettuce, Sports turf, Home lawns and Legume seed crops and pastures, as per the DIRECTIONS FOR USE TABLE

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

Contents: 500g, 1kg, 10kg, 20kg

APVMA Approval No: 68409/58305

**Sipcam Pacific Australia Pty Limited
Level 1, 191 Malop Street
Geelong VIC 3220**

STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. DO NOT re-use container or dispose of undiluted chemicals on-site. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation, and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling the dust. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier or from our web site, www.sipcam.com.au.

NOTICE TO BUYERS

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.

APVMA No.: 68409/58305

DOM:

Batch No:

This product is NOT considered to be a dangerous good under the Australian code for the transport of dangerous goods by road and rail

For specialist advice in emergency only,
call **1800 033 111** all hours, Australia wide

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Burst 500 WG Herbicide

ACTIVE CONSTITUENT: 500 g/kg PROPYZAMIDE

GROUP **D** HERBICIDE

For selective control of certain grasses and broad leaf weeds in Lettuce, Sports turf, Home lawns and Legume seed crops and pastures, as per the DIRECTIONS FOR USE TABLE

THIS LEAFLET FORMS PART OR THE LABEL

Contents: 500g, 1kg, 10kg, 20kg

APVMA Approval No: 68409/58305

Sipcam Pacific Australia Pty Limited
Level 1, 191 Malop Street
Geelong VIC 3220

DIRECTIONS FOR USE

Crops	Weeds Controlled	State	Rate kg / ha	Critical Comments	
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter grass, Rye grass, Summer Grass, Wireweed, Prince of Wales feather, Shepherd's Purse, Nettles.	All States	4.5 kg in 330 – 550L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
Transplanted				Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
		Vic only	2.3 kg in 300 – 550L	Apply on light sandy soils, low in organic matter	
Turf & lawns Common couch	Winter grass	All States	1.2 kg (12g/100m ²)	Rate/Bowling Green 37 m x 37 m 170g	Apply during the Autumn and Winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12 – 25 mm irrigation or rain and again 24 hour later. Repeat treatment if necessary on Qld blue couch after 4 weeks. DO NOT RESEED FOR 60 DAYS AFTER TREATMENT. Use ONLY on established turf. DO NOT use on bluegrass, ryegrass, fescue or bent grass lawns.
Queensland blue couch		Qld only	0.6 kg (6g/100m ²)	85g	
Buffalo grass		WA only	1.2 kg (12g/100m ²)	170g	
Lucerne, Clover, Medics etc. Grown for forage, Hay or Seed production	Annual Ryegrass, Winter Grass, Barley Grass, Silver grass, Prairie Grass, Canary Grass, Great Brome, Squirrel tail Fescue and Rats tail Fescue.	NSW, ACT, S A, WA, Tas only	1 – 1.5 kg	<p>SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed.</p> <p>ESTABLISHED CROPS: Apply any time before the end of July.</p> <p>WEEDS: Apply when weeds are at post or Pre-emergent stages.</p> <p>BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.</p>	
	Wild oats, Perennial Ryegrass, Bent grass, English couch, Yorkshire fog grass, Chickweed, Seedling Sorrel, Blackberry nightshade, Patterson's curse, seedling Wireweed.		1.5 – 2 kg		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION

NOT CLASSIFIED AS DANGEROUS GOODS FOR
TRANSPORT BY ROAD OR RAIL.

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

Spray Preparation

Burst WG Herbicide is a water dispersible granule formulation and must be thoroughly dispersed in water and applied as a dilute spray solution. Partly fill the boomspray-tank with water so to form a slurry. Add the required amount of Burst 500 WG Herbicide for the area to be sprayed into water containers. Add this correct amount of pre-mixed Burst 500 WG Herbicide into the boomspray-tank with the agitation system operating. Fill up to the correct volume with water. The mixture must be kept in suspension at all times by continuous agitation, otherwise bottom settling of the product will result. Re-dispersion of the product will be extremely difficult and blockages of filters are likely to occur. When mixing with other chemical products ensure to check all label instructions prior to chemicals being tank mixed with Burst 500 WG Herbicide to check for chemical compatibility. Test a small area for any adverse effects prior to application.

DO NOT mix with any products that contains the active constituent, MCPA. If tank mixing, Burst 500 WG Herbicide must be in suspension before adding other herbicides, insecticides or surfactant. Check all labels **Directions for Use** table prior to tank mixing with Burst 500 WG Herbicide. Wash sprayer thoroughly after use.

Equipment

Always ensure the sprayer is clean, using a suitable spray and boom cleaner as directed. It is recommended that for turf situations, up to 100 L/ha spray volume is required.

WEED RESISTANCE WARNING

GROUP D HERBICIDE

Burst 500 WG Herbicide is a member of the benzamide group of herbicides with inhibitors of microtubal assembly mode of action. For weed resistance management this is a Group D herbicide. Some naturally occurring weed biotypes resistant to Burst 500 WG Herbicide and other benzamide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Burst 500 WG Herbicide or other benzamide herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam accepts no liability for any losses that may result from the failure of Burst 500 WG Herbicide to control the resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a Sipcam representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

PROTECTION OF CROPS

DO NOT apply to, or allow spray to drift onto crops, pastures or non target – desirable plants, nor onto agricultural land scheduled for cropping or pasture, as injury to the crop or pasture may occur.

PROTECTION OF LIVESTOCK

DO NOT graze sheep, cut for stock feed (including poultry and pigs) any treated areas (for 25 days after application). DO NOT FEED TURF / LAWN CLIPPINGS TO POULTRY OR LIVESTOCK

STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. **DO NOT** re-use container or dispose of undiluted chemicals on-site. **DO NOT** dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation, and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling the dust. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier or from our web site, www.sipcam.com.au.

NOTICE TO BUYERS

This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.