READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Gro-Chem

Lime Sulphur

Fungicide and Insecticide

ACTIVE CONSTITUENT: 200 g/L Sulphur (S) as Polysulphide Sulphur

For the control of certain diseases, insects and mites on Citrus, Grapevines, Ornamentals, Pome and Stone fruits and Vegetables.

As specified in the Directions for Use Table.

Contents: 5 20 200 1000 Litre



APVMA Approval No. 68891/59314

DIRECTIONS FOR USE

Restraints:

DO NOT apply during the growing period of apricots and peaches or other sulphur sensitive plants.

DO NOT apply when freezing weather is predicted, or the air temperature is over 32°C. DO NOT apply within 2 weeks of an oil spray except as indicated under the compatibility heading.

Сгор	Pest/ Disease	State	Rate per 100L water	Critical Comments
CITRUS Grapefruit Oranges	White Louse Scale	NSW, ACT, Vic, SA & TAS only	10 L	Winter Spray: Apply in July and thoroughly wet all
Lemons Mandarins etc	Maori Mite	Qld only NSW, Vic,	3.0 - 6.5 L 2.5 L	parts of the tree. Summer Spray:
Triandamis ded	Maon Med	ACT, SA, WA & TAS only	2.5 L	Spray to thoroughly wet leaves.
GRAPEVINES	Bud Mite Erinose Mite	Vic, WA, SA & Tas only	10 L	Apply as near as possible to
	Rust Mite Vine Bunch Mite	Qld, ACT and NSW only	5 L	budburst. Note: Spread of mites is checked by sulphur treatments used for powdery mildew
	Post Harvest treatment of mites	SA only	2 L	As soon as possible after harvest.
POME FRUIT Apples	Black Spot (Scab)	All States	5 L	Apply during dormant to late bud swell. Note: Lime Sulphur used after late green tip will
	Mildew San Jose Scale	Qld, NSW, ACT, SA, Vic & WA only		restrict growth and depress yield. DO NOT use on Delicious or Cox's

Pears	Black Spot Byrobia mite Pear leaf blister mite	All states	5 L	Orange Pinnin apples. Apply during dormant to late bud swell. Note: Lime Sulphur used after late green tip will restrict growth and depress yield.
STONE FRUIT Almonds Apricots Peaches Plums Prunes Nectarines	Byrobia Mite Freckle Leaf Curl Rust Shot Hole Brown Rot (Blossom Blight) San Jose Scale Frosted Scale	All States Qld, NSW, ACT, SA, Vic & Tas only Qld, Vic, SA, WA & TAS only	5 L	Spray while trees are dormant to the bud swell stage.
ORNAMENTALS Hedges Roses VEGETABLES Tomatoes	Two spotted mite, Powdery mildew, Rust Tomato mite Two spotted mite, Powdery Mildew, Rust	All States	1 L	Apply as a Summer Spray. Spray to wet leaves and repeat as necessary.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Gro-Chem Lime Sulphur Fungicide Insecticide is readily miscible in water and should be used on the same day as it is mixed.

Mixing

Before using, shake container contents well. Add the required amount of Gro-Chem Lime Sulphur to the water in the spray tank with stirring or agitation. Agitate while spraying. DO NOT mix Gro-Chem Lime Sulphur with acidic water of pH below 4.

Application

Spray in the cool of the day. The spray mixture should be used on the same day it is mixed with water. DO NOT apply when freezing weather is predicted or the air temperature is over 32 degrees celsius.

DO NOT apply product within 2 weeks of an oil spray except as indicated under the Compatibility statement below.

Compatibility

Gro-Chem Lime Sulphur may be mixed with winter oil for use on pome and stone fruit during the dormant to semi-dormant stages only. For details consult the label of winter oil. Gro-Chem Lime Sulphur has a limited compatibility with other pesticides and the use of any mixture is entirely at the user's risk.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plant/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

draft label Lime Sulphur [AU] [submitted] 12 Mar 2013.docx

Avoid contact with the eyes and skin. When preparing spray and using the prepared spray wear elbow-length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For additional information, refer to MSDS available from your supplier.

CONDITIONS OF SALE

This high quality product is suitable for the purpose advised on the label but as Gro-Chem cannot exercise control over mixing or applications, all conditions and warranties, as to its use for any purpose, are excluded and no responsibility will be accepted by Gro-Chem for any damage or injury whatsoever arising from its storage or application.

In transport emergency dial 000. Police or Fire Brigade For specialist advice in an emergency only, call 1800 777 068, 24 Hours.

Registered to:

Batch no:

Distributed by:

Agrinova NZ Ltd

E.E. Muirs and Sons
18 Raymond Road

PO Box 50 080

Date of Manufacture:

Laverton North VIC 3026 Australia 03 99312200

15 Sunlight Grove Porirua

New Zealand

Phone:

1800 777 068

Email:

grochem@grochem.co.nz