

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**CONQUEST**  
**NEON 400**  
**HERBICIDE**

**ACTIVE CONSTITUENT: 400 g/L FLUROXYPYR**  
**present as the METHYLHEPTYL ESTER**

**SOLVENT: 316 g/L LIQUID HYDROCARBON**  
**100 g/L N-METHYL-2-PYRROLIDONE**



**RLP**  
**Approved**

**GROUP I HERBICIDE**

For the control of a wide range of broadleaf Weeds in Fallow, Lucerne, Maize, Millets, Pastures, Poppies, Sorghum, Sugar cane, Sweet corn, Winter Cereals. Also for the control of Woody Weeds in Agricultural Non-Crop areas, Commercial and Industrial Areas, Forests, Pastures and Rights-of-way, as specified in the Directions for Use.

**Important: Read the attached leaflet before use.**

**Conquest Crop Protection Pty Ltd.**  
ABN 84 098 814 932  
Level 1, 4 Collingwood Street  
Osborne Park, WA 6017

Telephone (08) 9347 0500  
Facsimile (08) 9347 0551



## **STORAGE AND DISPOSAL**

### **Storage for all containers**

Store in closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

### **Disposal**

#### **Recycled containers:**

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container before disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at a drumMUSTER collection or similar container management site. The cap should not be replaced but may be taken separately.

#### **Non-recycled containers:**

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

#### **Refillable containers**

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## **SMALL SPILL MANAGEMENT**

Wear protective equipment (See **SAFETY DIRECTIONS**). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see **STORAGE AND DISPOSAL** section).

If necessary wash the spill area with an alkali detergent and water and absorb the wash liquid for disposal.

## **SAFETY DIRECTIONS**

Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves, a face shield or goggles. Wash hands after use.

After each day's use, wash gloves, face shield or goggles and contaminated clothing.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26).

If swallowed DO NOT induce vomiting. Give a glass of water.

## **MATERIAL SAFETY DATA SHEET**

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

## **CONDITIONS OF SALE**

The use of Conquest Neon 400 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Conquest Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Conquest Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

APVMA Approval No: 68984/59563

DOM:

Batch No:

**For specialist advice in an emergency dial  
1 800 033 111  
24 hours Australia wide.**

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**CONQUEST**  
**NEON 400**  
**HERBICIDE**

**ACTIVE CONSTITUENT: 400 g/L FLUROXYPYR**  
**present as the METHYLHEPTYL ESTER**

**SOLVENT: 316 g/L LIQUID HYDROCARBON**  
**100g/L N-METHYL-2-PYRROLIDONE**

**GROUP I HERBICIDE**

For the control of a wide range of broadleaf Weeds in Fallow, Lucerne, Maize, Millets, Pastures, Poppies, Sorghum, Sugar cane, Sweet corn, Winter Cereals. Also for the control of Woody Weeds in Agricultural Non-Crop areas, Commercial and Industrial Areas, Forests, Pastures and Rights-of-way, as specified in the Directions for Use.

**THIS LEAFLET IS PART OF THE LABEL**

**CONTENTS: 1 – 1000 LITRES**

**Conquest Crop Protection Pty Ltd.**  
ABN 84 098 814 932  
Level 1, 4 Collingwood Street  
Osborne Park, WA 6017

Telephone (08) 9347 0500  
Facsimile (08) 9347 0551



## DIRECTIONS FOR USE

### RESTRAINTS

**DO NOT** apply to plants which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected) poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.

Thorough coverage of both foliage and stems, to the point of runoff, is essential for high volume applications (see **GENERAL INSTRUCTIONS**; application methods **WOODY WEED SITUATIONS** section).

**DO NOT** spray if rain is likely to occur within one hour.

Table 1 Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures.

Table 2 Established Grass Pastures (Ground and Aerial)

Table 3 Sorghum, Maize, Millets and Sweet Corn

Table 4 Winter Cereals (Wheat, Barley, Oats and Triticale)

Table 5 Summer Fallow

Table 6 Winter Fallow

Table 7 Sugar Cane

Table 8 Lucerne (established only)

Table 9 Poppies

**Table 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures.**

- Legumes present at the time of spraying will be severely damaged.

<b>HIGH VOLUME APPLICATION: Dilute product with water.</b>				
<b>See General Instructions – Application Method for application details</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE per 100L water</b>	<b>CRITICAL COMMENTS</b>
Bathurst burr Noogoora burr	Seedlings and young plants up to 40 cm high	NSW, NT, Qld, WA only	38 mL	Add Uptake* Spraying Oil (see <b>General Instructions</b> ; Oils and surfactants).  <b>Boom spray:</b> Neon 400 at 0.3 L/ha + 0.4 L/ha of 2,4-D amine (625 g/L)
Black bindweed (climbing buckwheat)	Seedlings and young plants before flowering	NSW, Qld only	150 mL	
<i>Mimosa pigra</i>	Apply from mid to late summer	NT, WA only		
Common sensitive plant	Seedlings and young plants up to flowering	Qld, WA only	250 mL	
Bellyache bush		Qld, NSW, WA only		
Blackberry nightshade		NSW, Qld only		
Bokhara clover				
Caltrop (yellow vine) ( <i>Tribulus terrestris</i> ) ( <i>T. micrococcus</i> )	Seedlings and young plants up to 30 cm diameter			
Cobblers pegs	Up to 15 cm high			
Cockspur thorn	Up to 3 m high			
Creeping lantana	At flowering			
Crofton weed Mistflower	Seedlings and young plants up to flowering			
Docks ( <i>Rumex</i> spp.)	Seedlings and rosettes up to 30 cm high			
Hexham scent	Seedlings and young plants up to flowering			
Honey locust	Seedlings and young plants up to 2 m high			
Small flowered mallow ( <i>Marshmallow</i> ) ( <i>Malva parviflora</i> )	Seedlings and young plants up to flowering			
Yellowflower Devil's claw	Seedlings and young plants up to flowering			

**Table 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures.**

HIGH VOLUME APPLICATION: Dilute product with water. See General Instructions – Application Method for application details				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE per 100 L of water	CRITICAL COMMENTS
Lantana	Seedlings and regrowth 0.5 to 1.2 m high	NSW, Qld only	250 mL	Apply to actively growing plants from October to April. Some regrowth may occur particularly when treating old woody plants with sparse canopies.
	Plants and regrowth 1.2 to 2 m high		500 mL	
Blue heliotrope	Flowering		250 mL	
Limebush	Infestations up to 1.5 m high only			
Madeira vine	Apply at time of active growth			
Milkweed ( <i>Euphorbia heterophylla</i> )	3 leaf to flowering	Qld only	500 mL	Repeat applications will be necessary to control subsequent germinations.
Common sowthistle	Seedlings and young plants up to bolting	NSW, Qld only	250 mL	Add a surfactant (see <b>GENERAL INSTRUCTIONS</b> ; Oils and surfactants).
Mother-of-millions ( <i>Kalanchoe</i> spp.)	Seedling and young plants before flowering		300 mL	
Prickly acacia	Seedling and young plants up to 2 m high	Qld only	375 mL	Add Uptake* Spraying Oil (see <b>GENERAL INSTRUCTIONS</b> ; Oils and surfactants). Consult Tropical Weeds Research Centre, Charters Towers, for specific advice on application
<i>Sida</i> spp.	Seedling and young plants up to flowering	NSW, NT, Qld, WA	500 mL	
Broadleaf Pepper tree ( <i>Schinus terebinthifolius</i> )	Mature leaves, fruiting	Qld only	250 mL	Winter application only. Contact Alan Fletcher Research Station for more information.
Flannel weed ( <i>Sida cordifolia</i> )				
Snakeweed (Dark and light blue)	Seedling and young plants before flowering		375 mL	Add Uptake* Spraying Oil (see <b>GENERAL INSTRUCTIONS</b> ; Oils and surfactants).
Stinking Passion Flower	Established plants and regrowth	Qld, NT, WA	225 mL	Use 70mL/15 L for a knapsack.
Wandering jew ( <i>Tradescantia albiflora</i> )	Young plants up to and including flowering	All States	750 mL	Some regrowth will usually occur and will require retreatment.
Wattles (including <i>Acacia aulacocarpa</i> , <i>A. decora</i> , <i>A. harpophylla</i> , <i>A. leiocalyx</i> , <i>A. salicina</i> )	Seedling plants or regrowth 0.5 to 1.2 m high	NSW, Qld only	250 mL	Apply to actively growing plants when soil moisture is plentiful. Some regrowth may occur particularly when treating old woody plants with sparse canopies and under dry conditions.
	Plants or regrowth 1.2 to 2.0 m high only		500 mL	

**Table 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures.**

<b>BASAL PARK AND CUT STUMP APPLICATION: Dilute product with diesel.</b>				
<b>See General Instructions – Application Method for application details</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE per 100L of DIESEL</b>	<b>CRITICAL COMMENTS</b>
Celtis ( <i>Celtis sinensis</i> )	<b>Basal Bark only:</b> Young plants up to 2 m high and 20 cm basal diameter	Qld only	1.8 L	Treat stems from ground level to where multi-stemmed trunks branch.
Chinee apple	Up to 15 cm basal diameter		1.5 L	With basal bark, treat circumference of stem to a height of 45cm from the ground. Contact the Land Protection Branch, Department of Lands, Qld, for further information on Chinee Apple.
Cockspur thorn	<b>Basal Bark only:</b> Up to 5 cm basal diameter		1 L	
Mimosa bush <i>Acacia farnesiana</i> )	Up to 5 cm basal diameter	Qld, WA only	1.5 L	
Prickly acacia	Up to 10 cm basal diameter	Qld only	750mL	
Honey locust	Plants up to 10 cm basal diameter	Qld, NSW only	750mL	With basal bark, treat circumference of stem to a height of 45cm from the ground. <b>For cut stump application</b> use a rate of 5L/100L diesel for all plant sizes. Contact the Land Protection Branch, Department of Lands, Qld, for further information on Honey Locust.
	Plants 10 to 20 cm basal diameter		1.5 L	
	Plants >20cm basal diameter		2.5 L	
Sisal hemp ( <i>Agave</i> spp.)	All growth stages	Qld only	1.5 L	Treat as an overall spray. Contact The Land Protection Branch, Department of Lands, Qld for advice to control large infestations.
			5 mL undiluted product per plant	Lever out centre of plant with crowbar and immediately treat the exposed cut area
<b>BROADCAST AND AERIAL APPLICATION: Dilute product with water.</b>				
<b>See General Instructions – Application Method for application details</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE per 100L water</b>	<b>CRITICAL COMMENTS</b>
<i>Mimosa pigra</i>	Actively growing plants	NT, WA only	1.5 L	Aerial application: Add Uptake Spraying Oil at the rate of 1 L/100 L spray mix. Apply to actively growing plants from mid to late summer. Contact the Department of Primary Industries and Fisheries, NT for further information.
<b>LOW VOLUME, HIGH CONCENTRATE APPLICATION: Use a drench gun or gas-powered gun.</b>				
<b>See General Instructions – Application Method for application details</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE per 100L water</b>	<b>CRITICAL COMMENTS</b>
Limebush	Isolated bushes up to 1.2 m high only	NSW, Qld only	500 mL	Apply a 50 mL dose per 5m <sup>2</sup> of bush surface area.
Tree violet ( <i>Hymenanthera dentata</i> )	Apply from late flowering to green fruit up to 1.2 m high	NSW only		Apply a 50 mL dose per cubic metre of bush

**Table 2: Established Grass Pastures**

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE per 100L water	CRITICAL COMMENTS
Blue billygoat weed. Common sensitive plant Giant sensitive plant Spinyhead sida	Apply before flowering	Qld, WA only	750 mL	Add Uptake Spraying Oil at 1 L/ha
St John's wort	Apply from bud to full bloom (usually late Nov to early Jan)	ACT, NSW and Vic only	1.5 L	Some regrowth will occur. Treat regrowth the following season for best results. Use at least 200 L water/ha.
Silverleaf nightshade	From onset of flowering to early berry-set (usually spring to mid-summer)	NSW only	375 mL or 190 mL + 1.2L -1.6L 2,4-D amine (625 g/L)	Add Uptake Spraying Oil at 1 L/ha. To ensure maximum effect, delay application until the majority of shoots have emerged. Follow-up treatment of regrowth is critical for best control.

**Table 3: Sorghum, Maize, Millets and Sweet corn (NSW & Qld only)**

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE per ha	CRITICAL COMMENTS
Sorghum	Apply when secondary roots are present, from 4 fully expanded leaves (15 cm tall) up to boot (also see CRITICAL COMMENTS)	Annual ground cherry Wild gooseberry ( <i>Physalis</i> spp.)	2 to 8 leaf Up to 15 cm tall	250 mL	<b>Sorghum:</b> From 8 leaf to boot stage, use dropper nozzles to prevent herbicide coming in contact with the crop's leaves and the growing point (meristem).
		Apple-of-Peru	15 to 30 cm tall	375 mL	
			Seedling plants up to 15 cm tall		
		Bathurst burr Noogoora burr	2 to 8 leaf Up to 20 cm tall	250 mL	
Maize & Sweet corn	Apply when secondary roots are present, from 3 fully expanded leaves (10 cm tall) up to just before tasselling (see CRITICAL COMMENTS)	Pigweed ( <i>Portulaca oleracea</i> )	20 to 50 cm tall	375 mL	<b>Maize and sweet corn:</b> From 6 leaf to just before tasselling, use dropper nozzles to prevent the herbicides coming in contact with the crop's leaves and the growing point (meristem).
			Up to 10 cm diameter	250 mL	
		10 to 30 cm diameter	375 mL		
Millets	Spray when secondary roots have developed, usually early to mid-tillering, and not later than before heads start to form at the base of tillers. (See CRITICAL COMMENTS)	Sesbania pea	2 to 6 leaf Up to 10 cm tall	750 mL	<b>Millets: DO NOT</b> use mixes with atrazine.  ( <sup>1</sup> ) This treatment may be slightly damaging to the crop. To minimise crop damage apply using dropper nozzles <b>at all crop stages.</b>
		Silverleaf nightshade (NSW only) ( <sup>1</sup> )	Full flower to early berry	375 mL + Uptake at 300mL/100L	
		Starburr ( <i>Acanthospermum hispidum</i> ) (Qld only)	Up to 12 leaf and before flowering	750 mL or 375 mL + 1.6 L atrazine (600 g/L)	
		Thornapples ( <i>Datura</i> spp.)	2 to 8 leaf Up to 15 cm tall	375 mL	
		Volunteer sunflower	2 to 5 leaf Up to 20 cm tall	500 mL	



Table 3: Sorghum, Maize, Millets and Sweet corn (NSW & Qld only)

Neon 400 in tank-mixes with atrazine: Sorghum, Maize and Sweet corn.					
CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE per ha	CRITICAL COMMENTS
Sorghum Maize & sweetcorn  (continued)	Spray when secondary roots have developed, usually early to mid-tillering and not later than before heads start to form at the base of the tillers (See CRITICAL COMMENTS)	<i>Amaranthus</i> spp. Including: Boggabri weed, Dwarf amaranth, Green amaranth, Redshank Anoda weed Bladder ketmia Black pigweed ( <i>Trianthema portulacastrum</i> ) Caltrope (yellow vine), including <i>Tribulus terrestris</i> , <i>T. microoccus</i> and <i>T. maximus</i> Cowvine (peach vine) ( <i>Ipomoea lonchophylla</i> ) Hairy wandering jew ( <i>Commelina benghalensis</i> ) Mintweed	Seedling plants up to 15 cm tall or rosettes up to 15 cm diameter	250mL + 1.2L of atrazine flowable (600g/L or 675g/L) or 375mL + 1.6L of atrazine flowable (600g/L or 1.1kg of atrazine 900g/kg granules)	<p><b>Use the low rate</b> (250mL + 1.2 L) when weeds are small (5-7 cm tall/ diameter).</p> <p><b>Use the high rate</b> (375mL + 1.6 L) when the weeds are larger (7 - 15 cm tall/ diameter).</p> <p>Neon 400 is generally more compatible with <b>Liquid atrazine products</b> (see <b>GENERAL INSTRUCTIONS; compatibility section</b>).</p> <p>Add a surfactant (See <b>GENERAL INSTRUCTIONS; Oils and surfactants</b>).</p> <p><b>DO NOT</b> add an oil to mixtures of Neon 400 and atrazine.</p>
		<i>Euphorbia davidii</i>	Cotyledons to 4 nodes up to 15 cm	500 mL + 1.6 L atrazine flowable (600 g/L)	
		Volunteer peanuts	Up to 15 cm diameter	500 mL + 3.7 L atrazine flowable (600 g/l)	
Sweet corn: Tasmania only					
CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE per ha	CRITICAL COMMENTS
Sweet corn only	3 to 5 leaf	Blackberry nightshade Volunteer potatoes	3 to 5 leaf	500 mL	

**Table 4: Winter Cereals (Wheat, Barley, Oats and Triticale)**

CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE per ha	CRITICAL COMMENTS
Apply from 3 leaf to flag (Zadoks 13 to 39)	Bedstraw ( <i>Galium tricornutum</i> )	3 to 6 whorl	Vic, SA, WA	500 mL <sup>(1)</sup>	<sup>(1)</sup> Add either Uptake or a surfactant (see GENERAL INSTRUCTIONS: Oils and surfactants).  Useful suppression only. <b>Mixtures:</b> Mixing partners with Neon 400 may reduce crop selectivity. Apply at crop growth sages according to the mixing partner's recommendation.  Plants 15 to 30 cm tall will only be suppressed.
	Cleavers ( <i>Galium aparine</i> )		NSW, Vic only		
	Black bindweed ( <i>Climbing buckwheat</i> )	2 to 4 leaf	NSW, Qld only	250 mL <sup>(1)</sup>	
		2 to 6 leaf		375 mL or 250 mL + 5 g Metsulfuron methyl <sup>(1)</sup>	
	Common sowthistle ( <i>Sonchus oleraceus</i> )	2 to 5 leaf		500mL	
	Deadnettle	2 to 6 leaf		750 mL or 250 mL + 5 g Metsulfuron methyl <sup>(1)</sup>	
	Spiny emex (Doublegee, Three cornered jack)	2 to 4 leaf	NSW, SA, Qld, WA		
	Prickly lettuce	2 to 5 leaf	NSW, Qld, Tas, Vic, WA	500 mL	
	Volunteer lupins	2 to 8 leaf	NSW, Vic, WA only	750 mL	
	Volunteer potato	10 to 15 cm tall	WA and Tas only		
	Wireweed	2 to 3 leaf	NSW, Qld, SA, Tas, Vic, WA		
			NSW and Qld only	250 mL + 5 g Metsulfuron methyl <sup>(1)</sup>	
Bittercress ( <i>Coronopus didymus</i> ) Mustards Shepherd's purse Turnip weed Wild radish Wild turnip	Up to 8 leaf and up to 15 cm diameter	Qld, NSW, Vic, SA, Tas, WA	250 mL to 1.5 L + Metsulfuron methyl <sup>(1)</sup> or Eclipse <sup>(1)</sup> or MCPA LVE or MCPA amine	The Neon 400 rate depends on what other weeds are present as listed above. See <b>Mixtures</b> comment above. Metsulfuron methyl (600g/kg) @ 5 g/ha (this mix does not control wild radish). Eclipse @ 5-7 g/ha (use the 5 g rate on turnip weed only). MCPA LVE (500 g/L) @ 700 mL/ha. MCPA Amine (500 g/L) @ 1.0 L/ha.	

**Table 5: Summer Fallow**

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE per ha	CRITICAL COMMENTS	
Annual ground cherry Wild gooseberry <i>Physalis</i> spp.)	2 to 8 leaf, up to 15 cm tall	NSW, Qld only	375 mL <sup>(2)</sup>	<p><sup>(1)</sup>Add Uptake* Spraying Oil (see <b>GENERAL INSTRUCTIONS</b>; Oils and surfactants).</p> <p>When mixing with Glyphosate 450 to control both grass and broadleaf weeds, refer to the Glyphosate 450 label for use rates and adjuvants recommended for the grasses (see <b>GENERAL INSTRUCTIONS</b>; compatibility section).</p> <p><sup>(2)</sup>Delay treatment until the maximum number of shoots have emerged, but before the onset of fruiting (late summer). DO NOT treat plants showing symptoms from previous treatment. Use the high rate when longer term weed control (6-10 months) is required and delay planting crops during this period. The low rate will require follow-up treatments.</p>	
Bathurst burr Noogoora burr	2 to 8 leaf, up to 20 cm tall	NSW, Qld, Vic, WA only			
Bellvine	Pre-flowering	NSW, Qld only	250 mL + 1.2 L Glyphosate 450		
Bladder ketmia	4 to 8 leaf, up to 10 cm tall				
Cowvine (Peach vine) <i>Ipomoea lonchophylla</i>	2 to 10 leaf up to 10 cm diameter	NSW, Qld only	250 mL + 1.0 L Glyphosate 450		
Caltrope (yellow vine), including <i>Tribulus terrestris</i> , <i>T. maximus</i> and <i>T. microccus</i>	Up to 15 cm diameter				
Pigweed <i>(Portulaca oleracea)</i>	Up to 10 cm diameter				375 mL <sup>(1)</sup>
	Up to 60 cm diameter				375 mL + 1.0 L Glyphosate 450
<i>Polymeria pusilla</i>	2 to 10 leaf up to 20 cm diameter				500mL <sup>(1)</sup> or 250mL + 1.2L Glyphosate 450
Rhynchosia	Seedlings to early flowering				500 mL <sup>(1)</sup> or 190 mL + 800 mL Glyphosate 450
Smallflower mallow or Marshmallow <i>(Malva parviflora)</i>	Up to 8 leaf up to 20 cm diameter				500 mL <sup>(1)</sup>
Thornapples <i>(Datura spp.)</i>	2 to 8 leaf up to 15 cm diameter	NSW, Qld, WA only	375 mL <sup>(1)</sup> or 250 mL + 1.2L Glyphosate 450		
Sesbania pea	2 to 6 leaf up to 10 cm tall	NSW Qld only	750 mL <sup>(1)</sup> or 250 mL + 1.2L Glyphosate 450		
Perennial Ground Cherry <i>(Physalis virginiana)<sup>(w)</sup></i>	Bud to early flowering up to 20 cm tall		750 mL of 1.5L <sup>(1)</sup>		
Silverleaf nightshade	Full flower to early berry-set (usually Dec – Feb)	NSW only	375 mL or 190 mL + 1.2 – 1.6 L 2,4-D amine (625 g/L)	Add Uptake Spraying Oil at the rate of 1 L/100 L spray mixture. To ensure maximum effect, delay application until the majority of shoots have emerged. Follow-up treatment will be required to control regrowth and is critical for optimum control. If wanting to prevent seed set repeat applications may be needed in the same season, although this does not lead to better long term control.	
Volunteer peanuts	Up to 15 cm diameter	Qld only	500 mL + 3.7 L atrazine flowable (600 g/L)	Add a surfactant (see <b>General Instructions</b> ; Oils and surfactants). <b>Important:</b> see <b>GENERAL INSTRUCTIONS</b> ; compatibility section).	
Volunteer sunflowers	2 to 5 leaf up to 20 cm	NSW, Qld only	500 mL	Add Uptake Spraying Oil (see <b>General Instructions</b> ; Oils and surfactants section).	