

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

ABASECT

INSECTICIDE / MITICIDE

Active constituents: 18 g/L ABAMECTIN
Solvent: 481 g/L LIQUID HYDROCARBON
Co-Solvent: 100 g/L N-METHYL-2-PYRROLIDONE



GROUP	6	INSECTICIDE
-------	---	-------------

For the control of pest mites on pears, apples, tomatoes, citrus, hops, strawberries and ornamentals. For the control of pest mites and native budworm on cotton. As specified in the Direction for Use table.

**IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY
BEFORE USING THIS PRODUCT.**

CONTENTS: 250mL, 1L, 5 – 1000 LITRES

APVMA Approval Number: 69001/59600

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Level 1, 4 Collingwood Street
Osborne Park, WA 6017

Telephone (08) 9347 0500
Facsimile (08) 9347 0551



GENERAL INSTRUCTIONS

Abasect Insecticide / Miticide is for the control of pest mites on pears, apples, tomatoes, citrus, hops, strawberries and ornamentals. For the control of pest mites and native budworm on cotton. Refer to accompanying leaflet for complete directions for use and product constraints.

INSECTICIDE RESISTANCE WARNING

GROUP **6** INSECTICIDE

For insecticide resistance management Abasect Insecticide / Miticide is a Group 6 insecticide. Some naturally occurring insect biotypes resistant to Abasect Insecticide / Miticide and other Group 6 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Abasect Insecticide / Miticide or other Group 6 insecticides are used repeatedly. The effectiveness of Abasect Insecticide/Miticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use Conquest Crop Protection Pty accepts no liability for any losses that may result from the failure of Abasect Insecticide / Miticide to control resistant insects. Abasect Insecticide / Miticide may be subject to specific resistant management strategies. For further information contact your local supplier, Conquest Crop Protection Pty representative or local agricultural department agronomist.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT use this container for any other purpose. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

For refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Conquest Crop Protection Pty .

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When opening the container, preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. For space applications wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves, goggles, half facepiece respirator with dust cartridge or canister. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with

soap and water. After each day's use, wash gloves, goggles, respirator and if rubber wash with detergent and warm water and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, DO NOT induce vomiting. Give a glass of water. **If skin contact occurs**, remove contaminated clothing and wash skin thoroughly. **If in eyes hold eyes open**, flood with water for at least 15 minutes and see a doctor.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for Abasect Insecticide / Miticide is available from Conquest Crop Protection Pty Ltd on request or via our website, www.conquestag.com.au

CONDITIONS OF SALE

Conquest Crop Protection Pty Ltd accepts no responsibility for the constituent quality of the product, however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions for use or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

UN No. 2902

PESTICIDES, LIQUID, TOXIC, N.O.S.*

(CONTAINS ABASECT)

PACKING GROUP III

HAZCHEM CODE 2XE



IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

EMERGENCY CONTACT

1 800 033 111

AUSTRALIA WIDE, 24 HOURS

Batch No.:

D.O.M.:

APVMA Approval Number: 69001/59600

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

ABASECT

INSECTICIDE / MITICIDE

Active constituents: 18 g/L ABAMECTIN
Solvent: 481 g/L LIQUID HYDROCARBON
Co-Solvent: 100 g/L N-METHYL-2-PYRROLIDONE

GROUP	6	INSECTICIDE
--------------	----------	--------------------

For the control of pest mites on pears, apples, tomatoes, citrus, hops, strawberries and ornamentals. For the control of pest mites and native budworm on cotton. As specified in the Direction for Use table.

**IMPORTANT: READ THE ATTACHED BOOKLET THOROUGHLY
BEFORE USING THIS PRODUCT.**

CONTENTS: 250mL, 1L, 5 – 1000 LITRES

APVMA Approval Number: 69001/59600

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Level 1, 4 Collingwood Street
Osborne Park, WA 6017

Telephone (08) 9347 0500
Facsimile (08) 9347 0551



DIRECTIONS FOR USE

RESTRAINTS:

DO NOT use if rainfall is expected before spray has dried as reduced efficacy may result.

DO NOT overhead irrigate within 24 hours after application.

SITUATION	PEST	RATE	WITHHOLDING PERIOD	CRITICAL COMMENTS
Cotton	Two Spotted Mite (<i>Tetranychus urticae</i>), Carminic mite (<i>Tetranychus cinnabarinus</i>)	300 mL/ha	20 days (harvest) 20 days (livestock/grazing)	Do NOT make more than two applications to cotton per season, regardless of pest being controlled. Apply by ground spraying equipment or fixed wing aircraft in a minimum volume of 20L/ha. Good coverage is essential. Monitor crops regularly and apply as soon as the threshold mite number for your area has been reached. Best results will be obtained when applied to low mite populations. Applications to high populations may not give satisfactory control. Under these conditions a second miticide application 7-10 days later may be needed. Do not use more than twice in one season for mite control. Refer to notes on resistance under general instructions section of this leaflet.
	Native budworm (<i>Helicoverpa punctigera</i>)	300 mL/ha OR 600 mL/ha		Use only when Lepton test kit results indicate no greater than 10% <i>Helicoverpa armigera</i> are present. Use the higher rate alone or the lower rate with a suitable mixing partner. Applications should be targeted at brown eggs and newly emerged larvae (neonates). Mixed size larval populations should be avoided. Do not use more than twice in one season for native budworm control. Do NOT make more than two applications to cotton per season, regardless of pest being controlled.
Apples, Pears	Two Spotted Mite (<i>Tetranychus urticae</i>), European Red Mite (<i>Panonychus ulmi</i>)	750 mL/ha plus 5 L/ha summer oil (See General Instructions for mixing rates.)	14 days (harvest) 14 days (livestock/grazing)	Apply as dilute or concentrated sprays but in no less than 1000L/ha. Good coverage is essential. Use a high quality summer spraying oil and follow label instructions. For best results apply soon after mite numbers have reached the threshold for your area. Generally Abasect Insecticide / Miticide takes about 7 days to reach maximum mite control. Abasect Insecticide / Miticide will control moderate to high mite populations but, in the absence of predatory mites (see integrated Pest Management below), re-treatment with another miticide may be necessary. If re-treatment is required, use an approved miticide from a different chemical group. Refer to notes on resistance and crop safety under General Instructions section of this leaflet. Integrated Pest Control: The effects of Abasect Insecticide / Miticide parasitic wasps and other beneficial insects in Australian orchards are not fully known. Studies have shown that after application of Abasect Insecticide / Miticide, predatory mite populations may not increase for a number of weeks, due to a lack of suitable pest mite prey. Predatory mite numbers will increase with any increase in pest mite numbers allowing the continuation of biological mite control. DO NOT use in IPM programs unless the pest mite threshold has been reached and predators are unlikely to achieve effective control.

DIRECTION FOR USE – Continues

SITUATION	PEST	RATE	WITHHOLDING PERIOD	CRITICAL COMMENTS
Citrus	Brown Citrus Rust Mite (<i>Tegolophus australis</i>), Citrus Rust Mite (<i>Phyllocoptera oleivora</i>), Broad Mite (<i>Polyhagotarsonemus latus</i>)	12 mL – 25 mL/100L plus 250 mL/100mL summer oil	7 days (harvest)	Apply as pest pressure indicates as a dilute spray in 3000 to 6000 L or water/ha. Use the higher rate under conditions of high pest pressure. Make no more than one application per season.
Hops	Two Spotted Mite (<i>Tetranychus urticae</i>)	1.0 L/ha	4 weeks (harvest/grazing/ cut for stockfood)	Apply as pest pressure indicates) as a dilute spray in 1000 to 2000 L Water/ha, depending on crop size. Make no more than one application per season.
Tomatoes - trellised or staked	Two Spotted Mite (<i>Tetranychus urticae</i>)	High volume spraying: 300 – 450 mL/ha Abasect Insecticide / Miticide using concentrations of 60 – 90 mL/100L	3 days (harvest) 3 days (livestock/ grazing)	Wet foliage to near the point of run-off. Thorough coverage and penetration into bushes is essential. Preferably apply before the build-up of mite numbers. If mite numbers exceed 5-6 mites per compound leaf, use the higher rate. Re-apply as pest numbers indicate. Allow at least 28 days between applications with Abasect Insecticide / Miticide and do not use more than two Abasect Insecticide / Miticide sprays per crop. Refer to notes on resistance under General Instructions section of this leaflet.
	Tomato Russet Mite (<i>Aculops lycopersici</i>)			Apply as for Two Spotted Mite. The lower rate will control Tomato Russet Mite not apparent at spraying. Use the higher rate when Tomato Russet Mite is present at spraying or is the main pest.
	Tobacco Leafminer (Potato moth) (<i>Phthorimaea operculella</i>)	High volume spraying: 600 mL/ha using concentrations of 120 mL/100L		Wet foliage to near the point of run off. Thorough coverage and penetration into bushes is essential. Apply on the first sign of pests. Re-apply as pest numbers indicate or every 7-10 days with a maximum of 5 applications to the crop. If mites are also a problem DO NOT use more than two Abasect Insecticide / Miticide sprays per crop. Refer to notes on Resistance under General Instructions section of this label. NB. Other IPM strategies to reduce leafminer populations should be employed.
Tomatoes - Field grown (not trellised)	Two Spotted Mite (<i>Tetranychus urticae</i>)	300 – 450 mL/ha	3 days (harvest) 3 days (livestock/ grazing)	Apply in sufficient volume to obtain even coverage and penetration of plants. Use droppers to direct spray onto plants and away from the inter-row. Preferably apply before the build-up of mite numbers. If mite numbers exceed 5 mites per compound leaf, use the higher rate. Re-apply as pest numbers indicate. Allow at least 28 days between applications with Abasect Insecticide / Miticide and DO NOT use more than two Abasect Insecticide / Miticide sprays per crop. Refer to notes on Resistance under General Instructions section of this leaflet.
	Tomato Russet Mite (<i>Aculops lycopersici</i>)			Apply as for Two Spotted Mite. The lower rate will control Tomato Russet Mite not apparent at spraying. Use the higher rate when Tomato Russet Mite is present at spraying or is the main pest.

	Tobacco Leafminer (Potato moth) (<i>Phthorimaea operculella</i>)	600 mL/ha		Apply in sufficient volume to obtain even coverage and penetration of plants. Use droppers to direct spray onto plants away from the inter-row. Apply on the first sign of pests. Re-apply as pest numbers indicate or every 7-10 days with a maximum of 5 applications to the crop. If mites are also a problem DO NOT use more than two Abasect Insecticide / Miticide sprays per crop. Refer to notes on Resistance under General Instructions section of this label. NB. Other IPM strategies to reduce leafminer populations should be employed.
Strawberries	Two Spotted Mite (<i>Tetranychus urticae</i>)	100 mL/100 L If spray volume is less than 600 L/ha, use a minimum of 600 mL Abasect Insecticide / Miticide per ha. Do not exceed 1200 mL/ha Abasect Insecticide / Miticide per Application.	3 days (harvest) 3 days (livestock/ grazing)	Spray to wet foliage to near the point of run-off. Thorough coverage and penetration into plants is essential. Preferably apply on first appearance of mites. When applied early, one application may be sufficient to give effective control however, if mite numbers exceed 3 - 5 mites per leaflet, apply two applications spaced 7-10 days apart. Re-apply as pest numbers indicate to a maximum of two sprays of Abasect Insecticide / Miticide per season. If retreatment is required after 2 consecutive sprays with Abasect Insecticide / Miticide, use an approved miticide from a different chemical group. Refer to notes on Resistance under General Instruction section of this leaflet. Integrated Pest Control: see comments in the Apple and Pear Section.
Ornamentals including Roses, Chrysanthemums, Carnations and indoor foliage plants	Two Spotted Mite (<i>Tetranychus urticae</i>)	50 mL/100 L to a maximum of 1.5 L Abasect Insecticide / Miticide per application.	DO NOT cut or graze for stock food	Spray to wet foliage to near the point of runoff using at least 2,000 litres per hectare (100 litres per 500 square metres). Thorough coverage and penetration into plants is essential. Preferably apply on first appearance of mites. If applied when pest numbers are low to moderate, one application will be sufficient to give effective control. However, if mites are numerous, apply two applications spaced 7-10 days apart. Re-apply as pest numbers indicate to a maximum of two sprays with Abasect Insecticide / Miticide per season. Ferns and Shasta Daisies may be affected by Abasect Insecticide / Miticide, so the product should not be used on them. For ornamentals not listed on the leaflet, small test applications to assess for unexpected phytotoxicity should be made before spraying the whole crop. DO NOT use more than two times per season. Refer to notes on Resistance under General Instructions section of this leaflet.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

COTTON: DO NOT HARVEST FOR 20 DAYS AFTER APPLICATION. DO NOT GRAZE CUT FOR STOCK FOOD FOR 20 DAYS AFTER APPLICATION.

APPLES, PEARS: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION. DO NOT FEED TREATED PRODUCE TO LIVESTOCK FOR 14 DAYS AFTER APPLICATION.

CITRUS: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

HOPS: DO NOT HARVEST, GRAZE OR CUT FOR STOCKFOOD FOR 4 WEEKS AFTER APPLICATION.

STRAWBERRIES, TOMATOES: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION. DO NOT FEED TREATED PRODUCE TO LIVESTOCK FOR 3 DAYS AFTER APPLICATION.

ORNAMENTALS: DO NOT CUT OR GRAZE FOR STOCK FOOD.

GENERAL INSTRUCTIONS

Abasect Insecticide / Miticide is for control of pest mites on pears, apples, tomatoes, citrus, hops, strawberries and ornamentals, and for the control of pest mites and native budworm on cotton. After application, Abasect Insecticide / Miticide quickly moves onto leaves where it remains for several weeks and where it is taken up by sucking mites. Abasect Insecticide / Miticide is not systemic, making good coverage essential. Abasect that is not absorbed into plants is quickly degraded.

INSECTICIDE RESISTANCE WARNING

GROUP	6	INSECTICIDE
--------------	----------	--------------------

For insecticide resistance management Abasect Insecticide / Miticide is a Group 6 insecticide. Some naturally occurring insect biotypes resistant to Abasect Insecticide / Miticide and other Group 6 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Abasect Insecticide / Miticide or other Group 6 insecticides are used repeatedly. The effectiveness of Abasect Insecticide / Miticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use Conquest Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of Abasect Insecticide / Miticide to control resistant insects. Abasect Insecticide / Miticide may be subject to specific resistant management strategies.

Therefore, Abasect Insecticide / Miticide should not be applied more than the following number of times per season/crop and should not be used consecutively except where a two-spray schedule is specifically recommended.

Apples, Pears, Citrus, Hops:	one spray per season
Tomatoes:	two sprays per crop if mites are present, or five sprays per crop if mites are not present
Cotton, Strawberries, Ornamentals:	two sprays per season

Alternate Abasect Insecticide / Miticide with approved miticides from other chemical groups. For further information contact your local supplier, Conquest Crop Protection Pty Ltd representative or local agricultural department agronomist.

CROP MONITORING

Effective control depends upon regular monitoring of crops. Check crops regularly (every 3-5 days) during the season.

APPLICATION ON COTTON

Air: Apply by air spray when there is a reliable cross wind to assist with good penetration into crop canopy. Application should be carried out in the cooler parts of the day or night. Preferably use aircraft fitted with micronair atomisers.

Ground: Use inter-row droppers fitted with nozzles spraying towards the cotton rows. Inter-row nozzles should be level with or just below the canopy and spraying at right angles to the ground.

MIXING

Measure the required amount of Abasect Insecticide / Miticide, add to the partly filled spray tank, and then add the remainder of the water. If oil is recommended, add this after the Abasect Insecticide / Miticide is well mixed. Wetting agent is not required.

MIXING INSTRUCTIONS FOR APPLES AND PEAR APPLICATIONS

To achieve 750mL Abasect Insecticide / Miticide plus 5L Summer Oil per ha, apply spray at the following mixing rates.

WATER	Abasect Insecticide / Miticide (Amount required per 100Litres)	Summer Oil
1000 L/ha (minimum)	75 mL	500 mL
1500 L/ha (minimum)	50 mL	335 mL
2000 L/ha (minimum)	37.5 mL	250 mL
2500 L/ha (minimum)	30 mL	200 mL

COMPATIBILITY

Abasect Insecticide / Miticide is compatible with most commonly used insecticides and fungicides. Do not apply with ULV formulations.

CROP SAFETY

Abasect Insecticide / Miticide plus Summer Oil may very occasionally cause slight russetting on some pear varieties, particularly ANJOU and other sensitive varieties when used alone or when other products are supplied sequentially. A very small amount of temporary apple fruit blemishing may be associated with low water volume applications. Abasect Insecticide / Miticide plus Summer Oil may cause more severe fruit russetting on sensitive apple varieties such as Golden Delicious when used within 7 days before or after application of the fungicide DELAN SC*. The Directions for Use and Precautions on summer oil labels should be followed carefully. Certain conditions may play a part in the occurrence of this damage. DO NOT make applications (1) when unusually hot conditions are present or expected within 24 hours after application, (2) under poor or slow drying conditions or (3) with equipment that may leave large droplets on fruit after application.

PRECAUTION

RE-ENTRY PERIOD: Under field conditions the spray should be allowed to dry on the foliage before re-entry into treated areas. DO NOT allow entry into treated areas in glasshouses for 24 hours after treatment. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and elbow-length PVC gloves.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do Not apply under weather conditions, or from spraying equipment, that may cause spray drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Studies indicate that when Abasect comes into contact with soil it readily and tightly binds to the soil and becomes inactive over time.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT use this container for any other purpose. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

For refillable containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Conquest Crop Protection Pty .

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When opening the container, preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. For space applications wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves, goggles, half facepiece respirator with dust cartridge or canister. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles, respirator and if rubber wash with detergent and warm water and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, DO NOT induce vomiting. Give a glass of water. **If skin contact occurs**, remove contaminated clothing and wash skin thoroughly. **If in eyes hold eyes open**, flood with water for at least 15 minutes and see a doctor.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for Abasect Insecticide / Miticide is available from Conquest Crop Protection Pty Ltd on request or through our website, www.conquestag.com.au.

CONDITIONS OF SALE

Conquest Crop Protection Pty Ltd accepts no responsibility for the constituent quality of the product, however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions for use or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

** not a Conquest Crop Protection trademark*

<p>UN No. 2902 PESTICIDES, LIQUID, TOXIC, N.O.S.* (CONTAINS ABASECT) PACKING GROUP III HAZCHEM CODE 2XE</p>
--

<p>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</p>
<p>EMERGENCY CONTACT 1 800 033 111 AUSTRALIA WIDE, 24 HOURS</p>