



Product Name: CONQUEST DARGO 500 WG SELECTIVE HERBICIDE
APVMA Approval No: 69009/107474

:

Label Name:	CONQUEST DARGO 500 WG SELECTIVE HERBICIDE
-------------	---

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	500 g/kg PROPYZAMIDE
-------------------------	----------------------

Mode of Action:	GROUP D HERBICIDE
-----------------	--------------------------

Statement of Claims:	For selective control of certain grasses and broad leaf weeds in canola, grain legumes, legume pasture, lettuce and turf according to the Directions for use Table
----------------------	--

Net Contents:	1kg-20kg 500g
---------------	------------------

Restraints:	Do not apply more than once per season Do not apply to water logged soils Do not apply to land where runoff to aquatic areas or sensitive crops like cereals is likely
-------------	--

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	
--------------------	--

Withholding Periods:	CANOLA: HARVEST – NOT REQUIRED WHEN USED AS DIRECTED
----------------------	--

	<p>GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION</p> <p>GRAIN LEGUMES: HARVEST – NOT REQUIRED WHEN USED AS DIRECTED GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION</p> <p>ALL OTHER CROPS: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION</p>
--	---

Trade Advice:	
---------------	--

General Instructions:	<p>GENERAL INSTRUCTIONS Best results are obtained by applying the product to cool, wet soil.</p> <p>Spray Preparation Dargo 500 WG Herbicide is a Wettable Granule formulation to be blended with water and applied as a spray. Granule formulations are designed to be added to large volumes of water and dispersed slowly under agitation. When using larger boomsprays and unable to add granules directly to a large volume in the tank via the top of the boomspray, pre-mixing may be necessary as follows:</p> <ul style="list-style-type: none"> • Partly (30%) fill the boomspray tank with water. • Slowly add required amount of Dargo 500 WG Herbicide for the area to be sprayed into Granni Pot® or other pre-mixing vessel while agitating well. • Ensure high water volumes and DO NOT add to the spray tank until the pre-mix is fully dissolved. This may take several mixes to ensure the correct amount is added to the boomspray tank. • Add the pre-mixed Dargo 500 WG Herbicide into the boomspray tank with the agitation system operating. • Fill up to the correct volume with water. • The mixture must be kept in suspension at all times by continuous agitation. • Tank-mixtures should be tested for physical compatibility and efficacy before use. <p>DO NOT mix with MCPA. If tank mixing, Dargo 500 WG Herbicide must be in suspension before adding other herbicides, insecticides or surfactants. Wash sprayer thoroughly after use. DO NOT handle the water-soluble packaging with wet hands or wet gloves. Keep dry.</p> <p>Equipment Always ensure the sprayer is clean, using a suitable spray and boom cleaner as directed. It is recommended that for turf situations, up to 100 L/ha spray volume is required.</p>
-----------------------	---

Resistance Warning:	<p>GROUP D HERBICIDE</p> <p>Conquest Dargo 500 WG Selective Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is a Group D herbicide. Some naturally occurring weed biotypes resistant to Conquest Dargo 500 WG Selective Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Conquest Dargo 500 WG Selective Herbicide or other amide herbicides.</p> <p>Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Pty Ltd accepts no liability for any losses that may result from the failure of Conquest Dargo 500 WG Selective Herbicide to control the resistant weeds.</p>
---------------------	--

	Do not rely exclusively on Dargo for weed control. Use as part of an integrated weed management program involving other modes of action and non-chemical methods of control. Advice as to strategies and alternative treatments that can be used, should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a Conquest' representative.
--	--

Precautions:	RE-ENTRY PERIOD Do NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.
--------------	--

Protections:	PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply to, or allow spray to drift onto crops, pastures or non target – desirable plants, nor onto agricultural land scheduled for cropping or pasture, as injury to the crop or pasture may occur. PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.
--------------	--

Storage and Disposal:	Store in the closed original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep away from contact with fertilizers, insecticides, fungicides and seeds. Protect from frost. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
-----------------------	---

Safety Directions:	Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.
--------------------	--

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
-------------------------	--

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE

Crops	Weeds Controlled	State	Rate kg/ha	Critical Comments	
Canola	Annual ryegrass, Winter grass, Barley grass, Silver grass, Prairie grass, Canary grass, Great brome, Squirrel tail fescue, Rats tail fescue	All States	1.0	Apply to moist soil prior to seeding only. Incorporate by sowing (IBS) with knife or blade points. A rainfall event (25 mm) between application and sowing, or soon after sowing is required for best results. Avoid seeding practices that moves treated soil back into the furrow. Heavy rainfall events that wash product back into the furrow can delay emergence and cause crop injury. Do not use on white sandy soils, or paddocks that have been recently mouldboard ploughed or spaded.	
Grain Legumes (including lupins, chickpeas, lentils, field peas, faba beans, navy beans)					
Legume Pastures (including lucerne, clover, medics etc. grown for forage, hay or seed production)					
	Wild oats, Perennial ryegrass, Bent grass, English couch, Yorkshire fog grass, Chickweed, Seedling sorrel, Blackberry nightshade, Patterson's curse, seedling wireweed	NSW, ACT, Tas, SA, WA only	1.0 - 1.5	Apply when weeds are at post or pre-emergent stages. SEEDLING CROPS: Apply after the third trifoliolate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July	
			1.5 – 2.0	Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.	
Lettuce (direct sown)	Barnyard grass, Blackberry nightshade, Chickweed, Portulaca, Winter grass, Rye grass, Summer grass, Wireweed, Prince of Wales feather, Shepherd's purse, Nettles	All States	4.5 kg in 330 – 550 L of water	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
Lettuce (transplanted)				Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
				Vic only	2.3 kg in 300 – 550 L of water
Turf and lawns Common couch	Winter grass	All States	1.2 kg (12 g/100 m ²)	Rate/Bowling Green 37 m x 37 m 170 g	Apply during the autumn and winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12 – 25 mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Qld blue couch after 4 weeks. DO NOT RESEED FOR 60 DAYS AFTER TREATMENT. Use ONLY on established turf. DO NOT use on bluegrass, ryegrass, fescue or bent grass lawns.
Queensland blue couch		Qld only	0.6 kg (6 g/100 m ²)	85 g	
Buffalo grass		WA only	1.2 kg (12 g/100 m ²)	170 g	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION