

Product Name: KENSO AGCARE KEN-MIDE 900 WG HERBICIDE  
APVMA Approval No: 69275 / 120864



Label Name:	KENSO AGCARE KEN-MIDE 900 WG HERBICIDE
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	900 g/kg PROPYZAMIDE
Mode of Action:	GROUP <b>D</b> HERBICIDE
Statement of Claims:	For selective control of certain grasses and broadleaf weeds in Canola, Legume Seed Crops and Pastures, Lettuce and Turf as per the Directions for Use table.
Net Contents:	250 g to 15 kg
Restraints:	
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	CANOLA: HARVEST: NOT REQUIRED WHEN USED AS DIRECTED. GRAZING: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 12 WEEKS AFTER APPLICATION.

ALL OTHER CROPS:  
DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS

Trade Advice:

General Instructions:

Best results are obtained by applying Kenso Agcare Ken-Mide 900 WG Herbicide to cool, wet soil.

#### MIXING

Ken-Mide 900 WG Herbicide is a Wettable Granule formulation to be blended with water and applied as a spray. Granule formulations are designed to be added to large volumes of water and dispersed slowly under agitation. When using larger boomsprays and unable to add granules directly to a large volume in the tank via the top of the boomspray. Pre-mixing may be necessary as follows:

- Partly (30%) fill the boomspray tank with water.
- Slowly add required amount of Ken-Mide 900 WG Herbicide for the area to be sprayed into pre-mixing vessel while agitating well.
- Ensure high water volumes and DO NOT add to the spray tank until the pre-mix is fully dissolved. This may take several mixes to ensure the correct amount is added to the boomspray tank.
- Add the pre-mixed Ken-Mide 900 WG Herbicide into boomspray tank with the agitation system operating.
- Fill up the correct volume with water.
- The mixture must be kept in suspension at all times by continuous agitation.
- Tank-mixtures should be tested for physical compatibility and efficacy before use.

DO NOT mix with MCPA, if tank mixing, Ken-Mide 900 WG Herbicide must be in suspension before adding other herbicides, insecticides or surfactants.

Wash sprayer thoroughly after use.

#### EQUIPMENT

Always ensure the sprayer is clean, using a suitable spray and boom cleaner as directed. It is recommended that for turf situations, up to 100L/ha spray volume is required.

Resistance Warning:

#### RESISTANT WEEDS WARNING GROUP D HERBICIDE

Kenso Agcare Ken-Mide 900 WG Herbicide is a member of the amide group of herbicides with inhibitors of microtubule assembly mode of action. For weed resistance management this is a Group D Herbicide. Some naturally occurring weed biotypes resistant to Kenso Agcare Ken-Mide 900 WG Herbicide and Group D Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Kenso Agcare Ken-Mide 900 WG Herbicide or Group D Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Kenso Agcare Australia Pty Ltd accepts no liability for any losses that may result from the failure of Kenso Agcare Ken-Mide 900 WG Herbicide to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or an Kenso Agcare representative.

Precautions:

Protections:

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT  
DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

	<p>DO NOT apply under weather conditions or from spraying equipment which may cause spray drift onto nearby susceptible plants, crops, cropping lands or pastures.</p> <p>PROTECTION OF LIVESTOCK  DO NOT graze sheep, cut for stock feed (including poultry and pigs) any treated areas (for 25 days after application).  DO NOT FEED TURF/LAWN CLIPPINGS TO POULTRY OR LIVESTOCK.</p>
--	---

Storage and Disposal:	<p>Store in the closed, original container in a cool, dry, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Keep from contact with fertilizers, insecticides, fungicides and seeds.</p> <p>DO NOT re-use container or dispose of undiluted chemicals on-site. DO NOT dispose of undiluted chemicals onsite.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation, and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or products.</p>
-----------------------	--

Safety Directions:	<p>Harmful if inhaled. Will irritate the eyes. Avoid contact with eyes and skin. Do not inhale dust. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves, face shield or goggles and disposable dust face mask covering mouth and nose. If products in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.</p>
--------------------	--

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

**DIRECTIONS FOR USE**

CROP	WEEDS CONTROLLED	STATE	RATE kg/ha	CRITICAL COMMENTS	
				Best results are obtained when applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 2 to 4 cm of soil within 24 hours of application. This is similar to 25mm rain.	
<b>BROADACRE CROPS</b>					
<b>Canola</b>	Annual Ryegrass, Barley Grass, Canary Grass, Great Brome, Prairie Grass, Rats Tail Fescue, Silver Grass, Squirrel Tail Fescue, Wild oats (canola), Winter Grass	All States	0.56	Incorporate by sowing (IBS) when weeds are at the pre-emergent stage.	
Legume Pastures  (including Lucerne, Clover, Medics etc. Grown for forage, hay or seed production)		NSW, ACT, TAS, SA, WA only	0.56-0.83	<b>SEEDLING CROPS:</b> Apply after the third trifoliolate leaf is fully developed. <b>ESTABLISHED CROPS:</b> Apply any time before the end of July. <b>WEEDS:</b> Apply when weeds are at post or pre-emergence stages.	
<b>VEGETABLES</b>					
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Rye Grass, Summer Grass,	All States	2.5 in 330-550L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
Lettuce (Transplanted)	Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles			Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
		VIC only	1.28 in 300-550L	Apply on light sandy soils, low in organic matter.	
<b>TURF AND LAWNS</b>				Rate/ Bowling Green 37x37m	
Common couch	Winter Grass	All States	0.7 (7g/100m <sup>2</sup> )	94g	Apply during the Autumn and Winter, if established turf is infested with seedling Winter Grass. Ensure uniform coverage. Irrigate immediately after application with 12-25mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Qld blue couch after 4 weeks. <b>DO NOT RESEED FOR 60 DAYS AFTER TREATMENT.</b> Use <b>ONLY</b> on established turf. <b>DO NOT</b> use on bluegrass, ryegrass, fescue or bent grass lawns.
Queensland blue couch		Qld only	0.3 (3g/100m <sup>2</sup> )	47g	
Buffalo grass		WA only	0.7 (7g/100m <sup>2</sup> )	94g	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**