



Product Name: CLIPSHIELD 60 SPRAY ON FOR SHEEP
APVMA Approval No: 69326/123655

Label Name:	CLIPSHIELD 60 SPRAY ON FOR SHEEP
Signal Headings:	FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	60 g/L CYROMAZINE
Claims:	When applied as directed will protect long wool sheep against fly strike for 11 weeks.
Net Contents:	20L 25L
Directions for Use:	READ DIRECTIONS FOR USE BEFORE USING THIS PRODUCT
Restrictions:	DO NOT USE on sheep which are producing, or may in the future produce milk that may be used or processed for human consumption. DO NOT USE on sheep with less than 6 weeks wool.
Contraindications:	Should not be used on open wounds e.g.: those resulting from marking or mulesing operations.
Precautions:	
Side Effects:	
Dosage and Administration:	This section contains file attachment. bytes

General Directions:	<p>SHAKE WELL BEFORE USE</p> <p>DO NOT DILUTE. Apply directly from the container using a suitable applicator gun. Except for small lambs under 15 kg, the dose volume is determined by the number of months since shearing. Apply CLIPSHIELD 60 SPRAY ON FOR SHEEP according to the diagram and tables in the Directions for Use.</p> <p>CLIPSHIELD 60 SPRAY ON FOR SHEEP may be applied to all breeds of sheep, providing protection against fly strike. CLIPSHIELD 60 SPRAY ON FOR SHEEP is applied as a low-volume treatment onto or near the area where protection is required. When applying, the nozzle of the applicator gun should be approximately 25 cm above the fleece, to achieve a 15cm wide band at each pass.</p>
Withholding Periods:	<p>MEAT: DO NOT USE less than 7 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE on sheep which are producing or may in the future produce milk that may be used or processed for human consumption.</p> <p>WOOLHARVEST INTERVAL: DO NOT USE less than 2 months before shearing or fibre collection.</p>
Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 28 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 1300 255 347 or the APVMA website (www.apvma.gov.au/residues/).</p> <p>Use of this product may result in wool residues that may not comply with European Union environment quality standards.</p> <p>Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.</p>
Safety Directions:	<p>Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.</p>
First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.</p>
First Aid Warnings:	
Additional User Safety:	<p>Additional information is listed in the Safety Data Sheet.</p> <p>Sheep Re-handling Interval: After treatment wait until sheep are dry before re-handling.</p>
Environmental Statements:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT</p> <p>DO NOT contaminate dams, rivers or streams with the product or used containers.</p>
Disposal:	<p>This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash</p>

outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

Storage:

Store below 30oC (Room Temperature). Store in original container tightly closed in a dry, cool place. Do not store in direct sunlight. Do not re-use the container. Keep out of reach of children.

DOSAGE AND ADMINISTRATION

MONTHS SINCE SHEARING	TOTAL VOLUME/SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NUMBER OF BANDS	NUMBER OF SHEEP TREATED PER 20L[25L]	CRITICAL COMMENTS
FOR THE PREVENTION OF BODY STRIKE					<ol style="list-style-type: none"> 1. Use only a Landmark recommended Spray-On applicator gun 2. Ensure the applicator gun is clean. 3. Calibrate applicator gun using an accurate volume measure. 4. Ensure all areas to be protected are treated. 5. When preventing body strike ensure bands overlap along the midline – see diagram. 6. This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die. 7. Should not be mixed with any other product. 8. Should not be applied during gusty or windy weather as spray drift will reduce the period of protection. 9. Heavy rain following application of this product could diminish the period of protection. 10. CLIPSHIELD 60 SPRAY ON FOR SHEEP is coloured for band identification. This scourable dye will disappear shortly after treatment
up to 6	34	17	2	588[735]	
7	38	19	2	524[657]	
8	44	22	2	452[568]	
9	50	25	2	400[500]	
10	56	28	2	356[446]	
Lambs less than 15 kg	17	17	1	1176[1470]	
FOR THE PREVENTION OF BODY & CRUTCH STRIKE					
up to 6	51	17	3	392[490]	
7	57	19	3	348[439]	
8	66	22	3	300[378]	
9	75	25	3	264[333]	
10	84	28	3	236[297]	
Lambs less than 15 kg	34	17	2	588[735]	
FOR THE PREVENTION OF POLL STRIKE					
Rams and sheep susceptible to poll strike should be treated. Apply at least 5 mL of CLIPSHIELD 60 SPRAY ON FOR SHEEP to the areas requiring protection, around the poll and horns.					

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.