

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

AC CAPRICE 570 EC

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 570g/L MCPA PRESENT AS THE 2-ETHYL HEXYL ESTER

| | | |
|-------|---|-----------|
| GROUP | 1 | HERBICIDE |
|-------|---|-----------|

For the selective control of certain weeds in agricultural crops as listed in the
Directions for Use table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT

CONTENTS: 20L, 110L, 200L, 1000L

AXICHEM PTY LIMITED

7 PALM AVENUE
SHORNCLIFFE QLD 4017
Ph: (07) 3269 7437 Fax: (07) 3869 2741



Batch No:
Date of Manufacture:

APVMA Approval No: 69443/60809

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers and product.

Refillable Containers: (1000L only):

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of sale for refill or storage.

Envirodrum 100L Minim Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Axichem Pty Limited.

SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes. Will irritate the skin. When preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow length PVC gloves and goggles. If product in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with the directions under normal conditions of use. No warranty of merchantability for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions.

IN CASE OF EMERGENCY: PHONE 000 POLICE OR FIRE BRIGADE.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

AC CAPRICE 570 EC

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 570g/L MCPA PRESENT AS THE 2-ETHYL HEXYL ESTER

| | | |
|-------|----------|-----------|
| GROUP | 1 | HERBICIDE |
|-------|----------|-----------|

For the selective control of certain weeds in agricultural crops as listed in the
Directions for Use table.

**IMPORTANT: READ THIS BOOKLET THOROUGHLY BEFORE USING THIS
PRODUCT**

AXICHEM PTY LIMITED

7 PALM AVENUE
SHORNCLIFFE QLD 4017
Ph: (07) 3269 7437 Fax: (07) 3869 2741

APVMA Approval No: 69443/60809

DIRECTIONS FOR USE:

| Situation & Crop | Weeds Controlled | Rate/Ha | Critical Comments |
|---|--|---|--|
| Wheat (up to early flag leaf (Z37) only) | Refer Weed Table | 440 mL – 1.84L | Use the lower rate (440mL/ha) when crop is at 3-5 leaf stage (Z13-Z15). Apply higher rates from 5 leaf stage through to flag leaf just visible (Z15-Z37). Kulin: Use only from Z15-Z37 (WA only). |
| Oats, Barley, Triticale and Cereal Rye (up to early flag leaf (Z37) only) | | | Use the lower rate (440mL/ha) when crop is at 3-5 leaf stage (Z13-Z15). Apply higher rates from 5 leaf stage through to flag leaf just visible (Z15-Z37). Barley: Use only from 5 leaf stage to flag leaf just visible (Z15-Z37). |
| Grass Pastures and Grass Seed Crops | | | Apply to established grass based pastures. Damage may occur to Legumes if present. See Legume Tolerance under Protection of Crops. NOTE: Capeweed, Paterson's Curse (Salvation Jane) and Variegated Thistle may be poisonous to stock after spraying. |
| CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye | Wild Radish Stage of Weed Growth | | # Reduced efficacy (suppression only) may be achieved on wild radish larger than 8 leaf or greater than 180mm in diameter. |
| | Up to the 4 leaf stage and not more than 120 mm in diameter | AC Caprice 570 EC Selective Herbicide 175 mL plus Nugrex* 350 mL | DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics. |
| | Up to the 6 leaf stage and not more than 150 mm in diameter. | AC Caprice 570 EC Selective Herbicide 175 mL plus Nugrex** 500 mL | Crop Stage: Nugrex* 350mL + AC Caprice 570 EC Selective Herbicide 175mL: Apply from 3 leaf stage to fully tillered (Z13-30) |
| | Up to the 8 leaf stage and not more than 180 mm in diameter# | AC Caprice 570 EC Selective Herbicide 350 mL plus Nugrex** 500 mL | Nugrex** 500mL + AC CAPTRICE 570 EC Selective Herbicide 175mL: Apply from 3 leaf stage to fully tillered (Z13-30). Nugrex** 500mL + AC CAPTRICE 570 EC Selective Herbicide 350mL: Apply from 5 leaf stage to fully tillered (Z16-30). Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post-sowing). WA ONLY: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). Warning: Nugrex may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "Crop Tolerance" Section of General Instructions on the Nugrex label). |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIODS:

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

GRAZING: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions. Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours following application. DO NOT spray weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

APPLICATION/ MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remainder of water.

BOOM SPRAYING – Use 30-120 litres of water per hectare.

AERIAL SPRAYING – Use 10-90 litres of water per hectare.

EQUIPMENT MAINTENANCE AND USE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with Spraymate Tank & Equipment Cleaner followed by several clear water rinses.

COMPATIBILITY/TANK MIXES

This product can be tank mixed with Metsulfuron Methyl (600g/L), Dicamba (500g/L), Bromoxynil, Flowable Diuron, Flowable Terbutryn, Glyphosate, diclofop methyl (500g/L), triasulfuron (750g/l), Roundup[®], Grasp[®], Lontrel[®]/Archer[®], Propon, Puma[®], Glean[®]/Siege[®], Nugrex/Tigrex[®]. The effectiveness of Imidan[®], Le Mat[®], Metasystox and Ravensdown Unidime 400 Herbicide may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of Spraymate LI-700 is recommended. When tank mixing add the required amount of compatible product when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying. When mixing with diclofop methyl add an approved wetter surfactant and do not exceed 700 mL of AC Caprice 570 EC Selective Herbicide per hectare. For consistent results, use a minimum of 80 L water per hectare by ground and 30 L by air, and if wild oats are present, use 2 L of diclofop methyl (500g/L) per hectare.

RESISTANT WEEDS WARNING

| | | |
|-------|---|-----------|
| GROUP | I | HERBICIDE |
|-------|---|-----------|

AC Caprice 570 EC Selective Herbicide is a member of the Phenoxys group of herbicides. AC Caprice 570 EC Selective Herbicide has the disrupters of plant cell growth mode of action. For weed resistance management AC Caprice 570 EC Selective Herbicide is a Group I Herbicide. Some naturally occurring weed biotypes resistant to AC Caprice 570 EC Selective Herbicide and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by AC Caprice 570 EC Selective Herbicide or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Axichem Pty Limited accepts no liability for any losses that may result from the failure of AC Caprice 570 EC Selective Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

DRIFT WARNING

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, vines, tomatoes, vegetables, fruit trees, lucerne, legume crops and pastures, oilseed crops or other susceptible crops or trees (eg. kurrajongs, Belahs, eucalypts).

LEGUME TOLERANCE

The tolerance of pasture legumes to this product can vary with species, variety, growing conditions, stage of growth and companion crop cover. While MCPA is preferred to 2,4-D Amine in most situations, this product will cause greater damage than 2,4-D amine formulations. Lucerne and medics should not be sprayed under any circumstances with this product, as severe damage will result.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers and product.

Refillable Containers: (1000L only):

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of sale for refill or storage.

Envirodrum 100L Minim Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Axichem Pty Limited.

SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes. Will irritate the skin. When preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow length PVC gloves and goggles. If product in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with the directions under normal conditions of use. No warranty of merchantability for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions.

IN CASE OF EMERGENCY: PHONE 000 POLICE OR FIRE BRIGADE.

WEED TABLE: Specific Weed Control Recommendations

Where weeds are to be sprayed in a crop or pasture, use only the rates given for the particular crop or situation indicated under the Directions for Use table. In most cases, this will give control, however some hard to kill weeds in an advance stage of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and there is no crop or pasture. Use at these rates in a crop or pasture other than for spot spraying may cause damage.

| WEEDS CONTROLLED | RATE/ha | CRITICAL COMMENTS |
|---|-----------------|---|
| Bathurst Burr (<i>Xanthium spinosum</i>) | 1.4 L | Moderately susceptible. Spray young seedlings. |
| Capeweed (<i>Arctotheca calendula</i>) | 1.49 L | Apply up to 2-4 leaf stage only when plants are no larger than 5 cm diameter. |
| Charlock (<i>Sinapis arvensis</i>) | 400mL – 1.4 L | Young rosettes very susceptible. |
| Dandelion (<i>Taraxacum officinale</i>) | 1.4 L | Apply in early rosette stage. |
| Fat Hen (<i>Chenopodium album</i>) | 880mL – 1.4 L | Tolerance increases with age. Use lower rate only on seedlings. |
| Fumitory (<i>Fumaria</i>) | 965 mL | Moderately susceptible. Spray up to first true leaf stage (Not Tas). |
| Hedge Mustards (<i>Sisymbrium</i> spp) | 490 – 880 mL | Apply at rosette stage. |
| Lincoln Weed (<i>Diploaxis tenuifolia</i>) | 615 mL | Apply up to rosette stage only. |
| London Rocket (<i>Sisymbrium irio</i>) | 965 mL | Spray at young stage. |
| Noogoora Burr (<i>Xanthium occidentale</i>) | 1.4 L | Tolerance increases with age. Spray young seedlings only. |
| Paterson's curse (<i>Echium plantagineum</i>) | 1.49 L | Apply at 2 to 4 leaf stage only. |
| Saffron Thistle (<i>Carthamus lanatus</i>) | 965 mL – 1.66 L | Spray young rosettes before leaves become spiny. |
| Scotch (Cotton) Thistle (<i>Onopordum acanthium</i>) | 615 mL – 1.49 L | Spray young rosettes. |
| Skeleton weed (<i>Chondrilla juncea</i>) | 965 mL – 1.4 L | Moderately susceptible. Spray young rosettes before aerial growth commences. |
| Slender Thistle (<i>Carduus tenuiflorus</i>) | 1.84 L | Moderately susceptible. Spray young rosettes. |
| Spear Thistle (<i>Cirsium vulgare</i>) | 1.31 – 1.84 L | Moderately susceptible. Spray young rosettes. |
| Stinkwort (<i>Dittrichia graveolens</i>) | 1.4 L | Spray very young before plants 15cm high. |
| Turnip weed (<i>Rapistrum rugosum</i>) | 615 – 965 mL | Spray up to rosette stage. Apply lower rates to seedlings |
| Volunteer Canola (<i>Brassica napus</i>) | 1.31 L | Apply to seedlings only. |
| Variigated Thistle (<i>Silybum marianum</i>) | 735 – 965 mL | Spray at pre-cabbage stage |
| Volunteer Safflower (<i>Carthamus tinctorius</i>) | 880 mL | Apply to seedlings only. |
| Volunteer Sunflower (<i>Helianthus annuus</i>) | 1.31 L | Apply to seedlings only. |
| Wild Radish (<i>Raphanus raphanistrum</i>) | 965 – 1.4 L | Apply up to 2 to 3 leaf stage only. |
| Wild Sage (<i>Salvia verbenaca</i>) | 965 mL | Spray young seedlings. |
| Wild Turnip (<i>Brassica</i> spp) | 440 mL – 1.31 L | Spray young rosettes. Lower rates for seedlings. |