

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

TRIMMER

750 WG HERBICIDE



ACTIVE CONSTITUENT : 750 g/kg TRIBENURON-METHYL

GROUP **B** HERBICIDE

For the control of certain broadleaf weeds in fallows and pre-crop situations
as per the DIRECTIONS FOR USE table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

NET CONTENTS: 500g, 1kg, 2kg, 5kg

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Level 1, 4 Collingwood Street
Osborne Park, WA 6017

Telephone: (08) 9347 0500
Facsimile: (08) 9347 0551



STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. **DO NOT** re-use container. Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When preparing spray wear elbow-length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for Conquest Trimmer 750 WG Herbicide is available from Conquest Crop Protection Pty Ltd on request or via our website, www.conquestag.com.au

CONDITIONS OF SALE

Conquest Crop Protection Pty Ltd accepts no responsibility for the constituent quality of the product, however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions for use or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

<p style="text-align: center;">In a Transport Emergency Dial 000 Police or Fire Brigade</p>
<p style="text-align: center;">SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 033 111 ALL HOURS AUSTRALIA WIDE</p>

APVMA Approval No. : 69729 / 61510

Batch Number :

Date of Manufacture :

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

TRIMMER

750 WG HERBICIDE

ACTIVE CONSTITUENT : 750 g/kg TRIBENURON-METHYL

GROUP **B** HERBICIDE

For the control of certain broadleaf weeds in fallows and pre-crop situations
as per the DIRECTIONS FOR USE table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

NET CONTENTS: 500g, 1kg, 2kg, 5kg

Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Level 1, 4 Collingwood Street
Osborne Park, WA 6017

Telephone: (08) 9347 0500
Facsimile: (08) 9347 0551



DIRECTIONS FOR USE RESTRAINTS

DO NOT apply to weeds that are not actively growing or stressed by any cause such as adverse weather conditions, drought and water logging. Broadleaf weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or only growth suppression. If heavy grazing has occurred allow regrowth before spraying.

Minimum volumes of prepared spray to be applied: Ground application – 50L/ha
Aerial application – 20L/ha

Use prepared spray suspensions within two days of preparation, otherwise significant breakdown may occur.
DO NOT apply if rainfall is expected within 2 hours.

Conquest Trimmer 750 WG Herbicide alone – NSW & Qld ONLY

SITUATION	WEEDS CONTROLLED	RATE g/ha	WEEDS GROWTH STAGE AT APPLICATION	CRITICAL COMMENTS	
FALLOW and PRECROP (refer to Crop Options section for minimum intervals between application and sowing)	Amaranthus/ Boggabri weed (<i>Amaranthus mitchelli</i>)	25	Apply up to the 10 leaf stage	As a herbicide resistance Countering measure, DO NOT use an ALS inhibitor herbicide in the crop following the use of Conquest Trimmer 750 WG Herbicide as a preceding fallow or pre-crop treatment.	
	Caltrop/ Yellowvine (<i>Tribulus terrestris</i>)				
	Common Sowthistle/ Milk Thistle (<i>Sonchus oleraceus</i>)				
	Deadnettle (<i>Lamium amplexicaule</i>)	25	Apply up to the 8 leaf stage		If weeds are at a more advanced growth stage and/or are present in high numbers (greater than 50 per m ²) use a Glyphosate mixture as specified in Conquest Trimmer 750 WG Herbicide and Glyphosate Tank mix table.
	Medics (Native) (<i>Medicago</i> spp.)	30	Apply up to 5cm diameter		
	New Zealand Spinach (<i>Tetragonia tetragonoidea</i>)	20	Apply up to the 10 leaf stage		
	Prickly Lettuce (<i>Lactuca serriola</i>)	30	Apply up to the 4 leaf stage		If weeds are at a more advanced growth stage and/or are present in high numbers (greater than 50 per m ²) use a Glyphosate mixture as specified in Tank mix table.
	Tumip Weed (<i>Rapistrum rugosum</i>)	20	Apply up to flowering		

Always add non-ionic surfactant (1000g/L – non buffering type) at 100 mL/100L (0.1% v/v) of final volume. When applying Conquest Trimmer 750 WG Herbicide alone by air in addition to a non-ionic surfactant add a mineral spray oil at 1L/100L (1% v/v) of final spray volume.

Conquest Trimmer 750 WG Herbicide + Glyphosate Tank mixes NSW & Qld Only

For the control of these weeds in addition to those in the Conquest Trimmer 750 WG Herbicide alone table.

SITUATION	WEEDS CONTROLLED	RATE g/ha	WEEDS GROWTH STAGE AT APPLICATION	CRITICAL COMMENTS
FALLOW and PRE- CROP (refer to Crop Options section for minimum intervals between application and sowing).	Black Bindweed (<i>Fallopia convolvulus</i>)	25 + Glyphosate (450g/L) 800 mL/ha	Apply up to the 10-leaf stage	For best control, apply to small actively growing weeds. Larger weeds are more difficult to control.
	Common Thornapple (<i>Datura stramonium</i>)	20 + Glyphosate (450g/L) 400 mL/ha	Apply up to the 8-leaf stage	
	Deadnettle (<i>Lamium amplexicaule</i>)	25 + Glyphosate (450g/L) 400 mL/ha	Apply up to the 10-leaf stage	
	Mintweed (<i>Salvia reflexa</i>)	15 + Glyphosate (450g/L) 600 mL/ha		
	Pigweed/ Portulaca (<i>Portulaca oleracea</i>)	20 + Glyphosate (450g/L) 600mL/ha		
	Prickly Lettuce (<i>Lactuca serriola</i>)		Apply up to the 6-leaf stage	

Always add non-ionic surfactant (1000 g/L – non buffering type) at 100 mL/100L (0.1% v/v) of final spray volume.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Best results are obtained when Conquest Trimmer 750 WG Herbicide is applied to young actively growing weeds. The rate will depend on weed spectrum at time of application. The degree of control and duration of effect are dependent on rate used, sensitivity and size of target weeds and environmental conditions at the time of and following application.

Conquest Trimmer 750 WG Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms of dying weeds (discolouration) may not be noticeable for 1 to 3 weeks after application depending on the environmental conditions and susceptibility. Warm, moist conditions following treatment promote the activity of Conquest Trimmer 750 WG Herbicide, while cold, dry conditions delay activity. Weeds hardened-off by cold weather and/or drought stress will be less susceptible.

Where recommended the addition of Glyphosate will enhance knockdown and control of weeds. It also helps minimise the risk of resistant weeds occurring.

RESISTANT WEEDS WARNING

GROUP	B	HERBICIDE
--------------	----------	------------------

Conquest Trimmer 750 WG Herbicide is a member of the Sulfonylurea group of herbicides. Conquest Trimmer 750 WG Herbicide has the Inhibitor of Acetolactate Synthase (ALS) mode of action. For weed resistance management Conquest Trimmer 750 WG Herbicide is a Group B Herbicide. Some naturally occurring weed biotypes resistant to Conquest Trimmer 750 WG Herbicide and other Group B Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Conquest Trimmer 750 WG Herbicide or other Group B Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Crop Protection accepts no liability for any losses that may result from failure of Conquest Trimmer 750 WG Herbicides to control resistant weeds.

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be taken to prevent seed set of these survivors.

DO NOT use Group B Herbicide against the same weed in the crop following the use of Conquest Trimmer 750 WG Herbicide alone either as a fallow or pre-crop treatment. This does NOT apply if the fallow or pre-crop use if Conquest Trimmer 750 WG Herbicide was tank mixed with Glyphosate.

Avoid the prolonged use of Group B Herbicide on the same weed population.

If the user suspects that an ALS inhibitor resistant weed is present, Conquest Trimmer 750 WG Herbicide or other Group B Herbicides recommended for the control of those weeds should not be used.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Conquest Crop Protection representative.

CROP SAFETY

DO NOT apply chlorsulfuron, metsulfuron-methyl or triasulfuron products in a tank mix with Conquest Trimmer 750 WG Herbicide as a fallow or pre-crop treatment.

SPRAY PREPARATION

Conquest Trimmer 750 WG Herbicide is a dry flowable formulation to be mixed with water and applied as a spray. Partially fill the spray tank with water. Measure the amount of Conquest Trimmer 750 WG Herbicide required for the area to be sprayed. Add the correct amount of Conquest Trimmer 750 WG Herbicide to the spray tank with the agitation system engaged. Top up to the correct volume with water. **THE MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION.**

In tank mixes, Conquest Trimmer 750 WG Herbicide must be in suspension before adding the companion herbicide or surfactant/wetting agent.

COMPATIBILITY

Conquest Trimmer 750 WG Herbicide is compatible with Glyphosate herbicides.

USE OF SURFACTANT/WETTING AGENT

Always add non-ionic surfactant (1000g/L – non-buffering type) at 100 mL/100L (0.1% v/v) of final spray volume. When applying Conquest Trimmer 750 WG Herbicide alone by air in addition to a non-ionic surfactant add a mineral spray oil at 1L/100L (1% v/v) of final spray volume.

GROUND SPRAYING

Use a boom spray properly calibrated to a constant speed and rate of delivery to ensure thorough coverage and a uniform spray pattern. Avoid overlapping and shut off spray booms while starting, turning, slowing or stopping. Apply a minimum of 50L prepared spray/ha.

AERIAL SPRAYING

Apply a minimum of 20L/ha water. Application in 30L/ha will improve performance reliability. Avoid spraying in still conditions and in winds likely to cause drift onto adjacent sensitive crops or fallow areas likely to be planted to these crops – see Crop Options section. Turn off spray boom whilst passing over creeks and dams.

DO NOT apply when a temperature air inversion is likely to occur.

SPRAYER CLEANUP

To avoid subsequent injury to crops immediately after spraying thoroughly remove all traces of Conquest Trimmer 750 WG Herbicide from mixing and spray equipment as follows:

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through boom and hoses using at least one-tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water adding household ammonia to give a 1% solution (equivalent ratios include 1/3 litre household ammonia product, containing 9.5% ammonia) for each 100 litres of tank volume. Agitate, and then flush the boom and hoses with the cleansing solution. Top up with water making sure the tank is completely full and allow standing for 15 minutes with agitation engaged. Again flush the boom and hoses and drain tank completely.
4. Nozzles and screens should be removed and cleaned separately with ammonia solution at the same concentration as used for the sprayer.
5. Rinse the tank with clean water and flush through boom and hoses using at least one-tenth of the spray tank volume. Drain tank completely.
6. **DO NOT** spray the tank rinsate onto sensitive crop or land intended for cropping with sensitive crop.

NOTE: IF not possible to drain the tank completely, step 3 must be repeated before going on to step 4.

CROP OPTIONS

Land previously treated with Conquest Trimmer 750 WG Herbicide may be sown to any of the specified crops after the interval indicated in the following table:

Minimum interval	Crops**
3 days	Barley, Oats, Wheat
# 7 or 21 days	Maize, Mungbeans, Sorghum, Soybeans, Sunflowers

NDTE: Conquest Trimmer 750 WG Herbicide is broken down in soil, primarily by chemical hydrolysis, but to a lesser degree by microbial degradation. Breakdown is fastest in warm, wet acid soils and slower in cold, dry alkaline soils. For the above summer crops, if minimum soil temperatures at planting depth are less than or equal to 15°C for three consecutive days, then plant back intervals should be extended to 21 days.

** As a herbicide resistance countering measure, **DO NOT** use a Group B Herbicide in any of these crops following the use of Conquest Trimmer 750 WG Herbicide alone as a preceding fallow or pre-crop treatment. Tolerance of other crops (growth through to maturity) should be determined on a small scale before resowing into larger areas.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGETED PLANTS INCLUDING AQUATIC PLANTS.

Injury to or loss of desirable trees, vegetation including aquatic plants may result from failure to observe the following:
DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT.

DO NOT contaminate streams, rivers, and waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK

The use pattern is such that a withholding period for harvest or grazing is not required. It is recommended however, not to graze treated areas for 2 to 3 days to ensure adequate control is achieved.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. **DO NOT** re-use container. Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When preparing spray wear elbow-length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for Conquest Trimmer 750 WG Herbicide is available from Conquest Crop Protection Pty Ltd on request or via our website, www.conquestag.com.au

CONDITIONS OF SALE

Conquest Crop Protection Pty Ltd accepts no responsibility for the constituent quality of the product, however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions for use or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

<p>In a Transport Emergency Dial 000 Police or Fire Brigade</p>
<p>SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 033 111 ALL HOURS AUSTRALIA WIDE</p>

APVMA Approval No. : 69729 / 61510

Batch Number :

Date of Manufacture :

