#### READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**ACCENSI** 

# Phospot 200 Systemic Fungicide



ACTIVE CONSTITUENT: 200 g/L PHOSPHOROUS ACID present as MONO-DI POTASSIUM PHOSPHITE

## Controls Root Rot, Collar Rot, Downy Mildew. For Citrus, Avocadoes, Grapes & Ornamentals

Lahel

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

Leaflet:

IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING

Contents: 200mL - 1L

ACCENSI Pty Ltd ABN 94 079 875 184 60-76 Potassium Street, Narangba Qld 4504 Ph: (07) 3897 2000

APVMA Approval No.: 69853 61888



Batch Number

Date of Manufactu

#### HOW TO USE: Shake well before use.

DO NOT spray when shade temperatures exceed 35°C.

DO NOT spray when soil is dry and plants are suffering from moisture stress.

Applying as a protective spray will give more effective contrtol. Ensure thorough leaf coverage.

CROP	DISEASE	RATE	HOW TO APPLY
Young or small citrus and recently transplanted trees	Phytopthora Root Rot, Collar Rot	5-10 mL/L	Use higher rate under higher disease risk conditions (eg. for trees in heavy clay soil). Apply each year as a preventative foliar spray, in late Winter prior to flowering and again in
Mature citrus (except container grown Mandarin trees)		100-160 mL in 12 L water per tree	Autumn when fruit are approaching maturity. Spray until leaves are just wet. Use 5-10 mL/L rate for container grown Mandarin trees. Removal of fruit from affected trees will assist reovery.
Ornamentals	Phytopthora Root Rot, Collar Rot, Crown Rot	5 <b>m</b> L/L	Apply at 4-6 weekly intervals as foliar spray when conditions favour disease. Certain plants may be sensitive to phosphorus acid sprays. Test on a small sample first if in doubt. DO NOT apply when plants are dormant or stressed.
Avocado	Phytopthora Root Rot	2.5-3 mL/L	Apply as a foliar spray to run-off. Use 7.5-10L per adult tree. Apply every 3 weeks as a curative spray, or every 5-6 weeks as a preventative spray.
Grapes	Downy Mildew	6 mL/L	Apply as a foliar spray to the point of run-off. Apply at first sign of disease.

### NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED.

#### HOW TO PREPARE

Mix the required rate with water.

**CAUTION:** DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

#### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, dry place out of reach of children. DO NOT store in direct sunlight.

Dispose of empty container by wrapping in paper, placing in a plastic bag and putting in garbage.

#### SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

#### FIRST All

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

#### **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE:** The use of ACCENSI Phospot 200 Systemic Fungicide being beyond the control of the manufacturer no warranty expressed or implied is given by ACCENSI Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and ACCENSI Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.