

Company Name: Product Name:

ABBEY LABORATORIES PTY LTD BANG SPRAY-ON SHEEP BLOWFLY TREATMENT

APVMA Approval No: 8002

80022/100051

Label Name:	BANG SPRAY-ON SHEEP BLOWFLY TREATMENT	
Signal Headings:	FOR ANIMAL TREATMENT ONLY	
	READ SAFETY DIRECTIONS BEFORE OPENING OR USING	
		5

Constituent Statements:	CYROMAZINE 60g/L	
----------------------------	------------------	--

Claims: WILL PROTECT LONG WOOL SHEEP AGAINST FLY STRIKE FOR ELEVEN (11) WEEKS WHEN USED AS DIRECTED.	
--	--

Net Contents:	1L, 5L, 10L, and 20L

Restraints:	DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

Contraindications:	This product should not be used on sheep with less than 6 weeks wool as a reduced period of protection may result. This product is contraindicated for use on open wounds, e.g. those resulting from marking or mulesing operations. This product should not be applied during gusty or windy weather as spray drift will reduce the period of protection.
--------------------	---

Precautions:	

Side Effects:	

Dosage and Administration:	Dosage and Administration instructions are uploaded as an attachment since they contain tables for the dose rate and frequency of doses.	
	This section contains file attachment.	

File Name:	BANG SPRAY ON Dosage and Administration 1Dec14.docx
File Size:	15734 bytes

General Directions:	READ DIRECTIONS FOR USE BEFORE OPENING OR USING THIS PRODUCT
	DO NOT DILUTE. DO NOT mix with any other product.
	Apply directly from the container using BANG SPRAY-ON recommended applicator gun.
	Except for small lambs under 15kg, the dose volume is determined by the number of months since shearing.
	Apply BANG SPRAY-ON according to the tables in the Directions for Use.
	BANG SPRAY-ON maybe applied to all breeds of sheep, providing protection against flystrike.
	BANG SPRAY-ON is applied as a low-volume treatment onto or near areas where protection is required. When applying, the nozzle of the applicator gun should be
	approximately 25 cm above the fleece, to achieve a 15 cm wide band at each pass - see diagram.

Withholidng Periods:	MEAT: DO NOT USE less than 7 days before slaughter for human consumption. MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption. WOOL HARVEST INTERVAL: DO NOT USE less than 2 months before shearing or fibre collection.
-------------------------	---

Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 28 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the distributor on (02) 6452 4020 or the APVMA website (www.apvma.gov.au/residues).
	Use of this product may result in wool residues that may not comply with European Union environmental quality standards.
	Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

Safety Directions:	Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.
----------------------------	--

First Aid Warnings:	

Additional User	Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling.
Safety:	Additional information is listed in the safety data sheet.

Environmental	ENVIRONMENTAL PROTECTION:
Statements:	DO NOT contaminate dams, rivers or streams with the product or used containers.

.

Disposal:	Triple or preferably pressure rinse containers and dispose of rinsate in a disposal pit as
	described below. DO NOT dispose of undiluted chemicals on-site. Break, crush or puncture
	and bury empty containers in a local authority landfill. If not available bury the containers

below 500 mm in a disposal pit specifically marked and set up for this purpose clear of
waterways, vegetation and roots. Empty containers and product should not be burnt.

Storage:	Store below 30C (room temperature). Store in original container tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container. Keep out of reach of children.
----------	---

Name and Address:	James Findlay
	ABBEY ANIMAL HEALTH PTY LTD 165 SHARP ST, COOMA NSW 2630 INFO LINE: 02 6452 4020

,

BANG SPRAY- ON SHEEP BLOWFLY TREATMENT

DOSAGE AND ADMINISTRATION:

MONTHS SINCE SHEARING	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	SHEEP TREATED PER 1L
	FOR THE PREV	ENTION OF BOI	DY STRIKE	
Up to 6	34	17	2	29
7	38	19	2	26
8	44	22	2	22
9	50	25	2	20
10	56	28	2	17
Lambs less than 15 kg	17	17	1	58
FOR T Up to 6	HE PREVENTION	N OF BODY AND	CRUTCH :	STRIKE
7	57	19	3	17
8	66	22	3	15
9	75	25	3	13
10	84	28	3	11
Lambs less than 15 kg	34	17	2	29

FOR THE PREVENTION OF POLL STRIKE: Rams and sheep susceptible to poll strike should be treated. Apply at least 5 mL of BANG SPRAY- ON to the areas requiring protection, around the poll and horns.

CRITICAL COMMENTS:

1. Use BANG SPRAY-ON recommended applicator gun.

2. Ensure the applicator gun is clean.

3. Calibrate applicator gun using an accurate volume measure.

4. Ensure all areas to be protected are treated.

5. When preventing body strike ensure bands overlap along the midline - see diagram.

6. This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die. Use a diazinon knockdown product as a dressing for existing strike.

7. Heavy rain following the application of this product could diminish the period of protection.

8. BANG SPRAY-ON is coloured for band identification. This scourable dye will disappear shortly after treatment.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPOPRIATE LEGISLATION.