



Company Name: AGRICULTURAL PRODUCT SERVICES PTY LTD

Product Name: Whitestar Flutriafol SC Fungicide

APVMA Approval No: 80450/100886

Label Name:	Whitestar Flutriafol SC Fungicide
--------------------	-----------------------------------

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
-------------------------	---

Constituent Statements:	500g/l Flutriafol
--------------------------------	-------------------

Mode of Action:	GROUP 3 FUNGICIDE
------------------------	--------------------------

Statement of Claims:	For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser as per the Directions for Use table
-----------------------------	---

Net Contents:	10 Litres (110L, 200L & 1000L)
----------------------	-----------------------------------

Restraints:	
--------------------	--

Directions for Use:	This section contains file attachment. File Name: DIRECTIONS FOR USE Whitestar Flutriafol 500.docx File Size: 16099 bytes
----------------------------	---

Other Limitations:	
---------------------------	--

Withholding Periods:	APPLIED IN-FURROW AS A FERTILISER TREATMENT CANOLA, WHEAT AND BARLEY - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING
-----------------------------	---

Trade Advice:	
General Instructions:	<p>Fertiliser Application Granulated Fertiliser.</p> <p>To determine the correct volume of flutriafol 500 SC required to be applied, refer to Critical Comments in the Directions for use table, which provides a guide based on a range of common fertilisers application rates.</p> <p>Ensure that treated fertiliser is dry before sowing with the crop.</p> <p>Treated fertiliser should be applied in a band 3-4 cm below or to the side of the seed in planting furrow. It should not be broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered.</p> <p>Do not mix seed with treated fertiliser and Whitestar Flutriafol Fungicide SC.</p>
Resistance Warning:	<p>FUNGICIDE RESISTANCE WARNING GROUP 3 FUNGICIDE</p> <p>Whitestar Flutriafol SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Whitestar Flutriafol SC Fungicide is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Whitestar Flutriafol SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Whitestar Flutriafol SC Fungicide or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.</p> <p>Since the occurrence of resistant fungi is difficult to detect prior to use, Agricultural Product Services Pty Ltd accepts no liability for any losses that may result from the failure of Whitestar Flutriafol SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.</p>
Precautions:	<p>Re-entry or Re-handling</p> <p>Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.</p>
Protections:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p>
Storage and Disposal:	<p>STORAGE AND DISPOSAL 10L</p> <p>Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.</p> <p>Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.</p> <p>STORAGE AND DISPOSAL 110L – 1000L</p> <p>Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.</p>

Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Agricultural Product Services Pty Ltd should be advised immediately.

This container is reusable and remains the property of Agricultural Product Services Pty Ltd. DO NOT rinse empty container. Empty container fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Agricultural Product Services Pty Ltd for cleaning, relabelling and refilling.

Safety Directions:

Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

First Aid Instructions:

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26

First Aid Warnings:

DIRECTIONS FOR USE: ALL STATES

Crop	Disease	Rate per hectare	WHP	Critical Comments																					
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	200mL	4 weeks Grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 200mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.																					
				<table border="0"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr> <td>60</td> <td>333*</td> </tr> <tr> <td>70</td> <td>285</td> </tr> <tr> <td>80</td> <td>250</td> </tr> <tr> <td>100</td> <td>200</td> </tr> <tr> <td>120</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)	60	333*	70	285	80	250	100	200	120	166									
Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																								
60	333*																								
70	285																								
80	250																								
100	200																								
120	166																								
Wheat	Stripe rust (<i>Puccinia striiformis</i>)	100 or 200mL		For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected. Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.																					
	Septoria tritici blotch (<i>S. tritici</i>) Take-all (<i>Geaumannomyces graminis</i>)	200mL		<table border="0"> <thead> <tr> <th rowspan="2">Fertiliser rate per hectare (kg)</th> <th colspan="2">Volume of product per 100kg of fertiliser (mL)</th> </tr> <tr> <th>100mL rate</th> <th>200mL rate</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>200</td> <td>400 *</td> </tr> <tr> <td>60</td> <td>166</td> <td>333 *</td> </tr> <tr> <td>70</td> <td>143</td> <td>285</td> </tr> <tr> <td>80</td> <td>125 †</td> <td>250</td> </tr> <tr> <td>100</td> <td>100 †</td> <td>200</td> </tr> <tr> <td>120</td> <td>88 †</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)		100mL rate	200mL rate	50	200	400 *	60	166	333 *	70	143	285	80	125 †	250	100	100 †	200	120
Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																								
	100mL rate	200mL rate																							
50	200	400 *																							
60	166	333 *																							
70	143	285																							
80	125 †	250																							
100	100 †	200																							
120	88 †	166																							
Barley	Powdery mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>)	100 or 200mL		<p>* Application volumes over 300mL per 100kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers.</p> <p>† Where application volume is low, product may be diluted with water to improve coverage.</p>																					