



**Company Name:** AXICHEM PTY LTD  
**Product Name:** AC Flout 250 SC Fungicide

**APVMA Approval No:** 80777/101615

<b>Label Name:</b>	AC Flout 250 SC Fungicide
--------------------	---------------------------

<b>Signal Headings:</b>	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
-------------------------	---

<b>Constituent Statements:</b>	ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL
--------------------------------	--

<b>Mode of Action:</b>	<table border="1"><tr><td>GROUP 3 FUNGICIDE</td></tr></table>	GROUP 3 FUNGICIDE
GROUP 3 FUNGICIDE		

<b>Statement of Claims:</b>	For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use Table.
-----------------------------	---

<b>Net Contents:</b>	1L - 1000L
----------------------	------------

<b>Restrains:</b>	
-------------------	--

<b>Directions for Use:</b>	This section contains file attachment. <b>File Name:</b> AC FLOUT 250 SC FUNGICIDE DIRECTIONS FOR USE.docx <b>File Size:</b> 18638 bytes
----------------------------	--

<b>Other Limitations:</b>	
---------------------------	--

<b>Withholding Periods:</b>	WITHHOLDING PERIODS: A. AC FLOUT 250 SC APPLIED IN-FURROW AS A FERTILISER TREATMENT CANOLA, WHEAT AND BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING. B. AC FLOUT 250 SC APPLIED AS A FOLIAR SPRAY BARLEY – DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION
-----------------------------	--

	<p>- DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 10 WEEKS OF TREATMENT. WHEAT - DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION - DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 7 WEEKS OF TREATMENT.</p>
--	--

<b>Trade Advice:</b>	
----------------------	--

<b>General Instructions:</b>	<p><b>GENERAL INSTRUCTIONS</b> Fertiliser application Ensure the treated fertiliser is dry before sowing the crop. Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. DO NOT mix seed with treated fertiliser. Foliar Applications Check the crop carefully and regularly. Spray only if there is sufficient time for the rust to develop to a level on the variety in question when it will cause significant yield loss.</p>
------------------------------	--

<b>Resistance Warning:</b>	<p>Fungicide Resistance Warning</p> <p><b>GROUP 3 FUNGICIDE</b></p> <p>AC Flout 250 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide.</p> <p>Some naturally occurring individual fungi resistant to AC Flout 250 SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by AC Flout 250 SC Fungicide or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Axichem Pty Ltd accepts no liability for any losses that may result from the failure of AC Flout 250 SC Fungicide to control resistant fungi.</p>
----------------------------	--

<b>Precautions:</b>	
---------------------	--

<b>Protections:</b>	<p>Protection of Wildlife, Fish, Crustaceans and Environment DO NOT contaminate streams, rivers or waterways with chemical or used containers.</p>
---------------------	--

<b>Storage and Disposal:</b>	<p>Storage and Disposal Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose,</p>
------------------------------	--

clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**Safety Directions:**

Harmful if swallowed. Harmful if inhaled. May irritate the nose, throat and skin. May irritate the eyes. Avoid inhaling spray mist. Avoid contact with eyes and skin.  
When opening the container and preparing spray, wear elbow-length chemical resistant gloves. After each days use, wash gloves and contaminated clothing.  
Wash hands after use.

**First Aid Instructions:**

First Aid  
If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 131126).

**First Aid Warnings:**

**DIRECTIONS FOR USE  
FOR ALL STATES  
A. APPLIED AS AN IN-FURROW TREATMENT**

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS																				
Canola	Blackleg ( <i>Leptosphaeria maculans</i> )	400mL	4 weeks (grazing)	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400mL of the product on the quantity of fertiliser required to be sown in the furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333								
Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																							
60	667*																							
70	571																							
80	500																							
100	400																							
120	333																							
Wheat	Stripe rust ( <i>Puccinia striiformis</i> )	200 or 400mL		For the control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected. Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400mL of the product on the quantity of fertiliser required to be sown in the furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>50</td><td>400</td><td>800*</td></tr> <tr><td>60</td><td>333</td><td>667*</td></tr> <tr><td>70</td><td>286</td><td>571</td></tr> <tr><td>80</td><td>250*</td><td>500</td></tr> <tr><td>100</td><td>200*</td><td>400</td></tr> <tr><td>120</td><td>167*</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)	50	400	800*	60	333	667*	70	286	571	80	250*	500	100	200*	400	120	167*	333
	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																						
50	400	800*																						
60	333	667*																						
70	286	571																						
80	250*	500																						
100	200*	400																						
120	167*	333																						
	Septoria tritici blotch ( <i>S. tritici</i> ) Take-All ( <i>Geaumannomyces graminis</i> )	400mL																						
Barley	Powdery Mildew ( <i>Erysiphe graminis</i> ) Scald ( <i>Rhynchosporium secalis</i> )	200 or 400mL																						

**B. APPLIED AS FOLIAR SPRAY**

Wheat	Stripe rust ( <i>Puccinia striiformis</i> )	250-500mL <b>Ground application:</b> Apply in 100-150L water plus 200mL of a 1000g/L non-ionic surfactant per 100L of spray solution.	7 weeks (harvest or grazing)	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with a high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
	Leaf rust ( <i>P. recondita</i> ) Septoria nodorum blotch ( <i>Septoria nodorum</i> ), Septoria tritici blotch ( <i>S. tritici</i> )	<b>Aerial application</b> Apply in a minimum of 15L water plus 200mL a 1000g/L non-ionic surfactant per 100L of spray solution.		
Barley	Powdery mildew ( <i>Erysiphe graminis</i> )	Apply in a minimum of 15L water plus 200mL a 1000g/L non-ionic surfactant per 100L of spray solution.	10 weeks (harvest or grazing)	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. In crops with a high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.**