

Product Name: SPARTAN SPRAY ON SHEEP BLOWFLY PROTECTION

APVMA Approval No: 80858/116741

Label Name:	SPARTAN SPRAY ON SHEEP BLOWFLY PROTECTION
Signal Headings:	FOR ANIMAL TREATMENT ONLY
	READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	60 g/L Cyromazine
Claims:	When applied as directed will protect long wool sheep against fly strike for 11 weeks.
Net Contents:	1L, 5L, 20L
Directions for Use:	
Restraints:	Do not use on sheep which are producing or may in the future produce milk or milk products for human consumption.
Contraindications:	This product should not be used on sheep with less than 6 weeks wool as a reduced period of protection may result. This product should not be used on open wounds e.g. those resulting from marking or mulesing operations. This product should not applied during gusty or windy weather as spray drift will reduce the period of protection.
Precautions:	
Side Effects:	
Dosage and Administration:	

General Directions:

Do not dilute. Do not mix with any other product. Apply directly from the container using ZAGRO AUSTRALIA recommended spray-on applicator gun. Except for small lambs under 15Kg, the dose volume is determined by the number of months shearing. Apply Trooper Spartan according to the diagrams and tables in the Directions for Use. Trooper Spartan Spray may be applied to all breeds of sheep, providing protection against flystrike. Trooper Spartan Spray is applied as a low-volume treatment onto or near areas where protection is required. When applying, the nozzle of the applicator gun should be approximately 25cm above the fleece, to achieve a 15cm wide band at each pass.

Withholding Periods:

Withholding periods: Meat: Do not use less than 7 days before slaughter for human consumption. Milk: DO not use on sheep which are producing or may in the future produce milk or milk products for human consumption. WOOL HARVEST INTERVAL: DO NOT use less than 2 months before shearing or fibre collection.

Trade Advice:

Export Slaughter Interval (ESI): Do not use less than 28 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Zagro Customer Service on 1800 144 884 or the APVMA website www.apvma.gov.au/residues/ESI.shtml.

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

Safety Directions:

Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.

First Aid Instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

First Aid Warnings:

Additional User Safety:

Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling. Material Safety Data Sheet: If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy phone Zagro Australia Customer Service on 1800 144 884

Environmental Statements:

Do not contaminate dams, rivers or streams with the product or used containers.

Disposal:

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

Storage:

Store below 30°C (room temperature). Store in original container tightly closed in a dry, cool place. Do not store in direct sunlight. Do not re-use the container. Keep out of reach of children.

SINCE SHEARING	TOTAL VOLUME / SHEEP (ML)	VOLUME PER TREATMENT BAND (ML)	NO. OF BANDS	SHEEP TREATED PER 1L	1. Use a ZAGRO
FOR THE PR	EVENTION OF BODY S	STRIKE	ı		AUSTRALIA recommende Spray-On applicator gun.
Up to 6	34	17	2	29	2.Ensure the applicator gu is clean.
7	38	19	2	26	3.Calibate applicator gu using an accurate volum measure.
8	44	22	2	22	4.Ensure all areas to be protected are treated. 5.When preventing body strike ensure bands overlap along the midline.
9	50	25	2	20	
10	56	28	2	17	
Lambs less	17	17	1	58	6.This product is not idea for treating existing strikes
than 15 kg					Treated larvae may take 3-4
	EVENTION OF BODY A	IND CRUTCH STRIKE	:		days to die. 7.Heavy rain following th application of this producould diminish the period
	EVENTION OF BODY A	ND CRUTCH STRIKE	3	19	days to die. 7.Heavy rain following th application of this produce.
FOR THE PR				19	days to die. 7.Heavy rain following th application of this producould diminish the period of protection.
FOR THE PR	51	17	3		days to die. 7.Heavy rain following th application of this producould diminish the period of protection. 8. SPARTAN SPRAY-ON is coloured for
FOR THE PRI Up to 6	51 57	17	3	17	days to die. 7.Heavy rain following th application of this producould diminish the period of protection. 8. SPARTAN SPRAY-ON is coloured for band identification. This scourable dye will disappeas shortly after treatment.
FOR THE PRI Up to 6 7 8	51 57 66	17 19 22	3 3 3	17	days to die. 7.Heavy rain following th application of this producould diminish the period of protection. 8. SPARTAN SPRAY-ON is coloured for band identification. This scourable dye will disappee

Dosage & administration:

> Rams and sheep susceptible to poll strike should be treated. Apply at least 5 mL of TROOPER SPARTAN SPRAY-ON to the areas requiring protection, around the poll and horns.

Not to be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.