

Product Name: Albaugh Fluropix 400 Herbicide
APVMA Approval No.: 81092/141581



| | |
|--------------------------------|--|
| Label Name: | Albaugh Fluropix 400 Herbicide |
| Signal Headings: | CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING |
| Constituent Statements: | ACTIVE CONSTITUENT: 400g/L FLUROXYPYR present as the METHYL HEPTYL ESTER SOLVENT: 300g/L LIQUID HYDROCARBON 100g/L N-METHYL PYRROLIDONE |
| Mode of Action: | GROUP 4 HERBICIDE |
| Statement of Claims: | For the control of a wide range of broadleaf Weeds in Fallow, Lucerne, Maize, Millets, Pastures, Poppies, Sorghum, Sugar cane, Sweet corn, Winter Cereals. Also for the control of Woody Weeds in Agricultural Non-Crop areas, Commercial and Industrial Areas, Forests, Pastures and Rights-of-way, as specified in the Directions for Use. |
| Net Contents: | 1L, 5L, 20L, 60L, 110L, 200L, 500L, 1000L |
| Restraints: | DO NOT apply to plants which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected) poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result. Thorough coverage of both foliage and stems, to the point of runoff, is essential for high volume applications (see GENERAL INSTRUCTIONS; application methods WOODY WEED SITUATIONS section). DO NOT spray if rain is likely to occur within one hour. |
| Directions for Use: | This section contains file attachment. |

| | |
|---------------------------|--|
| Other Limitations: | |
|---------------------------|--|

| | |
|-----------------------------|---|
| Withholding Periods: | <p>GRAZING CROPS AND PASTURES: DO NOT GRAZE FAILED CROPS AND TREATED PASTURES OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.</p> <p>HARVEST POPPIES: DO NOT SPRAY POPPIES LATER THAN 10 WEEKS BEFORE HARVEST.</p> <p>OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED</p> |
|-----------------------------|---|

| | |
|----------------------|--|
| Trade Advice: | |
|----------------------|--|

| | |
|------------------------------|--|
| General Instructions: | This section contains file attachment. |
|------------------------------|--|

| | |
|----------------------------|--|
| Resistance Warning: | <p>RESISTANT WEEDS WARNING</p> <p>Albaugh Fluropix 400 Herbicide is a member of the pyridine group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group 4 Herbicide. Some naturally-occurring weed biotypes resistant to the product can other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individual can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 4 Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Albaugh Asia Pacific Limited accepts no liability for any losses that may results from the failure of this product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or Albaugh Asia Pacific Limited.</p> |
|----------------------------|--|

| | |
|---------------------|--|
| Precautions: | |
|---------------------|--|

| | |
|---------------------|---|
| Protections: | <p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS</p> <p>Susceptible crops include but are not limited to clovers, cotton, fruit, hops, lupins, ornamentals, peas, pine tree, potatoes, navy beans, safflower, shade trees, soybeans, sunflower, tobacco, tomatoes, vegetables and vines. Albaugh Fluropix 400 Herbicide can be damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected by Albaugh Fluropix 400 herbicide and establish quickly after treatment. Transitory damage can occur on some species particularly those that spread by stolons such as cough grass (<i>Cynodon dactylon</i>), Kikuyu grass and carpet grass (<i>Axonopus</i> sp.). DO NOT allow spray to drift onto susceptible crops, shade trees and Pinus spp. DO NOT use under weather conditions or from spraying equipment which could cause spray to drift onto nearby susceptible plants.</p> <p>PROTECTIONS OF LIVESTOCK</p> <p>DO NOT graze or cut treated crops or plants for food except as specified under withholding periods. Poisonous plants may become more palatable after spraying. DO NOT allow stock to re-enter paddocks containing treated poisonous plants, until the plants have died down.</p> <p>PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT</p> <p>DO NOT contaminate streams, rivers of waterways with chemical or used containers.</p> |
|---------------------|---|

| | |
|------------------------------|---|
| Storage and Disposal: | <p>STORAGE AND DISPOSAL</p> <p>Storage for all containers</p> <p>Store in closed, original container in a cool, well ventilated area. Do not store for prolonged</p> |
|------------------------------|---|

periods in direct sunlight.

Disposal (1L, 5L, 20L)

Triple or preferably pressure rinse containers before disposal. Add rinsings to tank. Do not dispose of chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or territory government regulations. DO NOT burn empty containers or product.

Refillable Containers (60L, 200L, 500L, 1000L)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

110L Mini Bulk Returnable Container

Do not tamper with the Micro Matic valve or security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty container to the point of purchase. This drum remains the property of Albaugh Asia Pacific Limited.

Safety Directions:

SAFETY DIRECTIONS

Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves, face shield or goggles. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

First Aid Instructions:

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26). If swallowed, do not induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

First Aid Warnings:

DIRECTIONS FOR USE

TABLE 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures.

- Legumes present at the time of spraying will be severely damaged.

| HIGH VOLUME APPLICATION: Dilute product with water. See General Instructions – Application Method for application details | | | | |
|--|--|--------------------------|--------------------|--|
| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE mL/100L WATER | CRITICAL COMMENTS |
| Bathurst burr, Noogoora burr | Seedlings and young plants up to 40 cm high | NSW, NT, Qld, WA only | 38 | Add Uptake* Spraying Oil (see General Instructions; Oils and surfactants). |
| Black bindweed (Climbing buckwheat) | Seedlings and young plants before flowering | NSW, Qld only | 150 | |
| Mimosa pigra | Apply from mid to late summer | NT, WA only | | |
| Common sensitive plant | Seedlings and young plants up to flowering | Qld, WA only | 250 | |
| Bellyache bush | | Qld, NSW, WA only | | |
| Blackberry nightshade, Bokhara clover | | NSW, Qld only | | |
| Calltrop (yellow vine) (Tribulus terrestris) (T. micrococcus) | Seedlings and young plants up to 30cm diameter | | | |
| Cobblers pegs | Up to 15 cm high | | | |
| Cockspur thorn | Up to 3 m high | | | |
| Creeping lantana | At flowering | | | |
| Crofton weed, Mistflower | Seedlings and young plants up to flowering | | | |
| Docks (Rumex spp.) | Seedlings and rosettes up to 30 cm high | | | |

TABLE 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures (continued)

| HIGH VOLUME APPLICATION: Dilute product with water. See General Instructions – Application Method for application details | | | | |
|--|---|---------------|--------------------|--|
| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE mL/100L WATER | CRITICAL COMMENTS |
| Hexham scent | Seedlings and young plants up to flowering | NSW, Qld only | 250 | Boom spray: Albaugh Fluropix 400 Herbicide at 0.3 L/ha + 0.5 L/ha of 2,4-D amine (500 g/L) |
| Honey locust | Seedlings and young plants up to 2m high | | | |
| Small flowered mallow (Marshmallow) (Malva parviflora) | Seedlings and young plants up to flowering | | | |
| Yellowflower, Devil's claw | Seedlings and young plants up to flowering | | | |
| Lantana | Seedlings and regrowth 0.5 to 1.2 m high Plants and regrowth 1.2 to 2 m high | | 500 | |
| Blue heliotrope | Flowering | | | Apply to actively growing plants from October to April. Some regrowth may occur particularly when treating old woody plants with sparse canopies. |
| Limebush | Infestations up to 1.5 m high only | | | |
| Madeira vine | Apply at time of active growth | | 250 | |
| Milkweed (Euphorbia heterophylla) | 3 leaf to flowering | Qld only | 500 | Repeat applications will be necessary to control subsequent germinations. |
| Common sowthistle | Seedlings and young plants up to bolting | NSW, Qld only | 250 | Add a surfactant (see GENERAL INSTRUCTIONS; Oils and surfactants). |
| Mother-of-millions (Kalanchoe spp.) | Seedling and young plants before flowering | | 300 | |
| Prickly acacia | Seedling and young plants up to 2 m high | Qld only | 375 | Add Uptake* Spraying Oil (see GENERAL INSTRUCTIONS; Oils and surfactants). Consult Tropical Weeds Research Centre, Charters Towers, for specific advice on application |

TABLE 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures (continued)

| HIGH VOLUME APPLICATION: Dilute product with water. See General Instructions – Application Method for application details | | | | |
|--|---|-----------------------|--------------------|--|
| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE mL/100L WATER | CRITICAL COMMENTS |
| Sida spp. | Seedling and young plants up to flowering | NSW, NT, Qld, WA only | 500 | |
| Broadleaf Pepper tree (Schinus terebinthifolius) | Mature leaves, fruiting | Qld only | 250 | Winter application only. |
| Flannel weed (Sida cordifolia) | | | | |
| Snakeweed (Dark and light blue) | Seedling and young plants before flowering | | 375 | Add Uptake* Spraying Oil (see GENERAL INSTRUCTIONS; Oils and surfactants). |
| Stinking Passion Flower | Established plants and regrowth | Qld, NT, WA only | 225 | Use 70mL/15 L for a knapsack |
| Wandering jew (Tradescantia albiflora) | Young plants up to and including flowering | All States | 750 | Some regrowth will usually occur and will require retreatment. |
| Wattles (including Acacia aulacocarpa A. decora A. harpophylla A. leiocalyx A. salicina) | Seedling plants or regrowth 0.5 to 1.2 m high | NSW, Qld only | 250 | Apply to actively growing plants when soil moisture is plentiful. Some regrowth may occur particularly when treating old woody plants with sparse canopies and under dry conditions. |
| | Plants or regrowth 1.2 to 2.0 m high only | | 500 | |

TABLE 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures (continued)

| BASAL BARK AND CUT STUMP APPLICATION: Dilute product with diesel. See General Instructions – Application Method for application details | | | | |
|--|--|---------------|----------------------------------|---|
| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE L/100L DIESEL | CRITICAL COMMENTS |
| Celtis (Celtis sinensis) | Basal Bark only: Young plants up to 2m high and 20 cm basal diameter | Qld only | 1.8 | Treat stems from ground level to where multi-stemmed trunks branch. |
| Chinese apple | Up to 15 cm basal diameter | | 1.5 | With basal bark, treat circumference of stem to a height of 45cm from the ground. Contact the Land Protection Branch, Department of Lands, Qld, for further information on Chinese Apple. |
| Cockspur thorn | Basal Bark only: Up to 5 cm basal diameter | | 1 | |
| Mimosa bush (Acacia farnesiana) | Up to 5 cm basal diameter | Qld, WA only | 1.5 | |
| Prickly acacia | Up to 10 cm basal diameter | Qld only | 0.75 | |
| Honey locust | Plants up to 10 cm basal diameter | Qld, NSW only | 0.75 | With basal bark, treat circumference of stem to a height of 45cm from the ground. For cut stump application use a rate of 5L/100 diesel for all plant sizes. Contact the Land Protection Branch, Department of Lands, Qld, for further information on Honey Locust. |
| | Plants 10 to 20 cm basal diameter | | 1.5 | |
| | Plants >20cm basal diameter | | 2.5 | |
| Sisal hemp (Agave spp.) | All growth stages | Qld only | 1.5 | Treat as an overall spray. Contact The Land Protection Branch, Department of Lands, Qld for advice to control large infestations. |
| | | | 5 mL undiluted product per plant | Lever out centre of plant with cut bar and immediately treat the exposed cut area. |

TABLE 1: Woody Weeds in Agricultural Non-Crop Areas and Rights-of-Way, Commercial and Industrial Areas, Forests and Pastures (continued)

| BROADCAST AND AERIAL APPLICATION: Dilute product with water. See General Instructions – Application Method for application details | | | | |
|---|-------------------------|-------------|-----------------|--|
| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE L/ha WATER | CRITICAL COMMENTS |
| Mimosa pigra | Actively growing plants | NT, WA only | 1.5 | Aerial application: Add Uptake Spraying Oil at the rate of 1 L/100L spray mix. Apply to actively growing plants from mid to late summer. Contact the Department of Primary Industries and Fisheries, NT for further information. |

| LOW VOLUME, HIGH CONCENTRATE APPLICATION: Use a drench gun or gas-powered gun. See General Instructions – Application Method for application details | | | | |
|---|---|---------------|-------------------|--|
| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE mL/10L WATER | CRITICAL COMMENTS |
| Limebush | Isolated bushes up to 1.2 m high only | NSW, Qld only | 500 | Apply a 50 mL dose per 5m ² of bush surface area. |
| Tree violet (Hymenantha dentata) | Apply from late flowering to green fruit up to 1.2 m high | NSW only | | Apply a 50 mL dose per cubic metre of bush. |

TABLE 2: Established Grass Pastures

| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE L/ Ha | CRITICAL COMMENTS |
|--|---|-----------------------|---|---|
| Blue billygoat weed, Common sensitive plant, Giant sensitive plant, Spinyhead sida | Apply before flowering | Qld, WA only | 0.75 | Add Uptake Spraying Oil at 1 L/ha |
| St John's wort | Apply from bud to full bloom (usually late Nov to early Jan) | ACT, NSW and Vic only | 1.5 | Some regrowth will occur. Treat regrowth the following season for best results. Use at least 200 L water/ha. |
| Silverleaf nightshade | From onset of flowering to early berry-set (usually spring to mid-summer) | NSW only | 375mL or 190mL + 1.2-1.6L 2,4-D amine 625 | Add Uptake Spraying Oil at 1 L/ha. To ensure maximum effect, delay the application until the majority of shoots have emerged. Follow-up treatment of regrowth is critical for best control. |

TABLE 3: Sorghum, Maize, Millets and Sweet Corn (NSW and Qld only)

| CROP | CROP GROWTH STAGE | WEEDS CONTROLLED | WEED GROWTH STAGE | RATE mL/ha | CRITICAL COMMENTS |
|----------------------|--|---|------------------------------------|--------------------------------------|--|
| Sorghum | Apply when secondary roots are present, from 4 fully expanded leaves (15 cm tall) up to boot (also see CRITICAL COMMENTS) | Annual ground Cherry, Wild gooseberry (Physalis spp.) | 2 to 8 leaf Up to 15 cm tall | 250 | Sorghum: From 8 leaf to boot stage, use dropper nozzles to prevent herbicide coming in contact with the crop's leaves and the growing point (meristem). Maize and Sweet Corn: From 6 leaf to just before tasselling, use dropper nozzles to prevent the herbicides coming in contact with the crop's leaves and the growing point (meristem). Millets: DO NOT use mixes with atrazine. (1) This treatment may be slightly damaging to the crop. To minimise crop damage apply using dropper nozzles at all crop stages. |
| | | Apple-of-Peru | 15 to 30 cm tall | 375 | |
| | | Bathurst burr, Noogoora burr | 2 to 8 leaf Up to 20 cm tall | 250 | |
| Maize and Sweet corn | Apply when secondary roots are present, from 3 fully expanded leaves (10 cm tall) up to just before tasselling (see CRITICAL COMMENTS) | Pigweed (Portulaca oleracea) | 20 to 50 cm tall | 375 | |
| | | | Up to 10 cm diameter | 250 | |
| Millets | Spray when secondary roots have developed, usually early to mid-tillering, and not later | | 10 to 30 cm diameter | 375 | |
| | | Sesbania pea | 2 to 6 leaf Up to 10 cm tall | 750 | |
| | | Silverleaf nightshade (NSW only) (1) | Full flower to early berry | 375 + LI 700 at 300mL/100L | |
| | | Starburr (Acanthospermum hispidum) (Qld only) | Up to 12 leaf and before flowering | 750 or 375 + 1.6L atrazine (600 g/L) | |
| | | Thornapples (Datura spp.) | 2 to 8 leaf Up to 15 cm tall. | 375 | |
| Volunteer Sunflower | 2 to 5 leaf Up to 20 cm tall. | 500 | | | |

TABLE 3: Sorghum, Maize, Millets and Sweet Corn (NSW and Qld only)

| AC Dozer Herbicide in tank-mixes with atrazine: Sorghum, Maize and Sweet corn. | | | | | |
|--|--|---|--|--|--|
| CROP | CROP GROWTH STAGE | WEEDS CONTROLLED | WEED GROWTH STAGE | RATE L/Ha | CRITICAL COMMENTS |
| Sorghum Maize and Sweet corn | Spray when secondary roots have developed, usually early to mid-tillering and not later than before heads start to form at the base of the tillers (See CRITICAL COMMENTS) | Amaranthus spp. including: Boggabri weed, Dwarf amaranth, Green amaranth, Redshank, Anoda weed, Bladder ketmia, Black pigweed (Trianthema portulacastrum), Caltrope (yellow vine), including Tribulus terrestris, T. microoccus and T. maximus, Cowvine (peach vine) (Ipomoea lanchophylla), Hairy wandering jew (Commelina benghalensis), Mintweed | Seedling plants up to 15 cm tall or rosettes up to 15cm diameter | 250mL + 1.2L of atrazine flowable (600 g/L) or 375mL + 1.6L of atrazine flowable (600 g/L) | Use the low rate (0.5 + 1.5L) when weeds are small (5-7 cm tall/ diameter). |
| | | | | | Use the high rate (0.75 + 2L) when the weeds are larger (7 - 15 cm tall/ diameter). |
| | | | | | Albaugh Fluropix 400 Herbicide is generally more compatible with Liquid atrazine products (see GENERAL INSTRUCTIONS; compatibility section). |
| | | | | | Add a surfactant (See GENERAL INSTRUCTIONS; Oils and surfactants). |
| | | | | | DO NOT add an oil to mixtures of Albaugh Fluropix 400 Herbicide and atrazine. |
| | | Euphorbia davidii | Cotyledons to 4 nodes up to 15 cm | 500mL + 1.6L atrazine flowable (600 g/L) | |
| | | Volunteer peanuts | Up to 15cm diameter | 500mL + 3.7L atrazine flowable (600 g/L) | |

Sweet Corn: Tasmania only

| CROP | CROP GROWTH STAGE | WEEDS CONTROLLED | WEED GROWTH STAGE | RATE L/Ha | CRITICAL COMMENTS |
|-----------------|-------------------|---|-------------------|-----------|-------------------|
| Sweet Corn only | 3 to 5 leaf | Blackberry nightshade, Volunteer potatoes | 3 to 5 leaf | 500mL | |

TABLE 4: Winter Cereals (Wheat, Barley, Oats and Triticale)

| CROP GROWTH STAGE | WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE mL/Ha | CRITICAL COMMENTS | |
|---|--|---------------------------------------|---------------------------------|---|--|--|
| Apply from 3 leaf to flag (Zadoks 13 to 39) | Bedstraw (Galium tricornutum) | 1 to 3 whorl | Vic, SA, WA only | 500 | (1) Add either Uptake or a surfactant (see GENERAL INSTRUCTIONS: Oils and surfactants). | |
| | Cleavers (Galium aparine) | | NSW, Vic only | | | |
| | Black bindweed (Climbing buckwheat) | 2 to 4 leaf | NSW, Qld only | 250 (1) | Useful suppression only. | |
| | | 2 to 6 leaf | | | | |
| | Common sowthistle (Sonchus oleraceus) | 2 to 5 leaf | | 500 | Mixtures: Mixing partners with Albaugh Fluropix 400 Herbicide may reduce crop selectivity. Apply at crop growth stages according to the mixing partner's recommendation. | |
| | Deadnettle | 2 to 6 leaf | | 750 or 250 + 5 g Metsulfuron methyl (1) | | |
| | Spiny emex (Doublegee, Three cornered jack) | 2 to 4 leaf | NSW, SA, Qld, WA only | | 750 | |
| | Prickly lettuce | 2 to 5 leaf | NSW, Qld, Tas, Vic, WA only | | 500 | |
| | Volunteer lupins | 2 to 8 leaf | NSW, Vic, WA only | | 750 | Plants 15 to 30 cm tall will only be suppressed. |
| | Volunteer potato | 10 to 15 cm tall | WA and Tas only | | | |
| | Wireweed | 2 to 3 leaf | NSW, Qld, SA, Tas, Vic, WA only | | 250 + 5 g Metsulfuron methyl (1) | |
| | | | NSW and Qld only | | | |
| | Bittercress (Coronopus didymus), Mustards, Shepherd's purse, Turnip weed, Wild radish, Wild turnip | Up to 8 leaf and up to 15 cm diameter | Qld, NSW, Vic, SA, Tas, WA only | | 250 to 750 + Metsulfuron methyl (1) or Eclipse (1) or MCPA LVE or MCPA amine | The Albaugh Fluropix 400 Herbicide rate depends on what other weeds are present as listed above. See Mixtures: comment above. Metsulfuron methyl (600 g/kg) @ 5 g/ha (this mix does not control wild radish). Eclipse @ 5 -7 g/ha (use the 5 g rate on turnip weed only). MCPA LVE (500 g/L) @ 700 mL/ha. MCPA Amine (500 g/L) @ 1.0 L/ha. |

TABLE 5: Summer Fallow

| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE mL/Ha | CRITICAL COMMENTS |
|--|---|------------------------|-------------------------------------|--|
| Annual ground cherry, Wild gooseberry (<i>Physalis</i> spp.) | 2 to 8 leaf, up to 15 cm tall | NSW, Qld only | 375 | (1) Add Uptake* Spraying Oil (see GENERAL INSTRUCTIONS; Oils and surfactants). |
| Bathurst burr Noogoora burr | 2 to 8 leaf, up to 20 cm tall | NSW, Qld, Vic, WA only | 250 + 1.2L Glyphosate 450 | When mixing with Glyphosate 450 to control both grass and broadleaf weeds, refer to the Glyphosate 450 label for use rates and adjuvants recommended for the grasses (see GENERAL INSTRUCTIONS; compatibility section). (2) Delay treatment until the maximum number of shoots have emerged, but before the onset of fruiting (late summer). DO NOT treat plants showing symptoms from previous treatment. Use the high rate when longer term weed control (6-10 months) is required and delay planting crops during this period. The low rate will require follow-up treatments. |
| Bellvine | Pre-flowering | NSW, Qld only | | |
| Bladder ketmia | 4 to 8 leaf, up to 10 cm tall | | | |
| Cowvine (Peach vine) (<i>Ipomoea lonchophylla</i>) | 2 to 10 leaf up to 10 cm diameter | | | |
| Caltrope (yellow vine), including <i>Tribulus terrestris</i> , <i>T. maximus</i> and <i>T. microccus</i> | Up to 15 cm diameter | | | |
| Pigweed (<i>Portulaca oleracea</i>) | Up to 10 cm diameter | | | |
| | Up to 60 cm diameter | | | |
| Polymeria pusilla | 2 to 10 leaf up to 20 cm diameter | | | |
| Rhynchosia | Seedlings to early flowering | | | |
| Smallflower mallow or Marshmallow (<i>Malva parviflora</i>) | Up to 8 leaf up to 20 cm diameter | | | |
| Thornapples (<i>Datura</i> spp.) | 2 to 8 leaf up to 15 cm diameter | NSW, Qld, WA only | 375 or 250 + 1.2L Glyphosate 450 | |
| Sesbania pea | 2 to 6 leaf up to 10 cm tall | | 750 or 250 + 1.2L Glyphosate 450 | |
| Perennial Ground Cherry (<i>Physalis virginiana</i>)(w) | Bud to early flowering up to 20 cm tall | NSW Qld only | 750 or 1.5L | |

TABLE 5: Summer Fallow (continued)

| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE mL/Ha | CRITICAL COMMENTS |
|-----------------------|--|---------------|--|--|
| Silverleaf nightshade | Full flower to early berry-set (usually Dec – Feb) | NSW only | 375 or 190 + 1.2L – 1.6L 2,4-D amine (500 g/L) | Add Uptake Spraying Oil at the rate of 1 L/100 L spray mixture. To ensure maximum effect, delay application until the majority of shoots have emerged. Follow-up treatment will be required to control regrowth and is critical for optimum control. If wanting to prevent seed set repeat applications may be needed in the same season, although this does not lead to better long term control. |
| Volunteer peanuts | Up to 15 cm diameter | Qld only | 500 + 3.7L atrazine flowable (600 g/L) | Add a surfactant (see GENERAL INSTRUCTIONS; Oils and surfactants). Important: see GENERAL INSTRUCTIONS; compatibility section). |
| Volunteer sunflowers | 2 to 5 leaf up to 20 cm | NSW, Qld only | 500mL | Add Uptake Spraying Oil (see GENERAL INSTRUCTIONS; Oils and surfactants section). |

TABLE 6: Winter Fallow

| WEEDS CONTROLLED | WEED GROWTH STAGE | STATE | RATE mL/Ha | CRITICAL COMMENTS |
|--|----------------------------------|------------------|---|---|
| Bedstraw (<i>Galium tricornutum</i>) | Up to 5 whorl | Vic, SA, WA only | 500 | 1) Add Uptake Spraying Oil (see GENERAL INSTRUCTIONS; Oils and surfactants section). (2) Add Uptake or a surfactant (see GENERAL INSTRUCTIONS; Oils and surfactants section). |
| Cleavers (<i>Galium aparine</i>) | | NSW, Vic only | | |
| Black bindweed (Climbing buckwheat) | 2 to 8 leaf up to 10 cm diameter | | 375 | When mixing with Glyphosate 450 to control both grass and broadleaf weeds, refer to the Glyphosate 450 label for use rates and adjuvants recommended for the grasses (see GENERAL INSTRUCTIONS; Compatibility Section). |
| Common sowthistle (<i>Sonchus oleraceus</i>) | 2 to 5 leaf up to 10 cm diameter | | 500 or 250 + 600mL Glyphosate 450 | |
| Prickly lettuce | | | | |
| Spiny emex (Doublegee, Three cornered jack) | 2 to 8 leaf | | 750 or 250 + 5g Metsulfuron methyl (600 g/kg) | |
| Wireweed | 2 to 3 leaf up to 10 cm tall | NSW, Qld only | 750 or 250 (2) + 5g Metsulfuron methyl (600 g/kg) or 500 + 600mL Glyphosate 450 | |

TABLE 7: Sugar Cane (Qld, NSW, NT and WA only)

| CROP GROWTH STAGE | WEEDS CONTROLLED | WEED GROWTH STAGE | RATE mL/ha | CRITICAL COMMENTS |
|----------------------------------|---|--|---|---|
| From early tillering to maturity | Balsum pear, Blackberry nightshade, Blue billygoat weed, Centro, Cowpea, Giant sensitive plant, Lablab bean, Noogoora burr, Phasey bean, Pinkburr, Prickly African Cucumber, Spinyhead sida, Stinking passion flower (seedlings only) | Apply from 2 to 3 leaf until flowering | Ground: 650 Aerial: 750 | For optimal weed control, delay application until just before the "close-in" stage. Aerial application: Apply in not less than 60 L/ha water and add Uptake Spraying Oil at 1L/100L spray mixture. Ground application: Apply in 100 – 400 L/ha water and add Uptake Spraying Oil at 500mL/100L of spray mixture |
| | Belvine, Morning glory, Red or pink convolvulus, Star-of-Bethlehem | | As above + 800mL 2,4-D amine 625 | |
| | Stinking passion flower | Established or ratoon plants with at least 1.0 m of regrowth | High volume: 225 mL/100 L water Knapsack: 35 mL/15 L water | Thoroughly wet plants to the point of run-off. |
| | Milkweed (Euphorbia heterophylla) | Seedlings and young plants up to flowering. | 1.5L or 1.15L + 3.3L atrazine flowable (600 g/L) | Better control will be achieved with the atrazine mixture. Delay application until just before the cane reaches the "close-in" stage. This will improve control and minimise the number of seedlings that germinate. |

TABLE 9: Poppies (Tas only)

| CROP STAGE GROWTH | WEEDS CONTROLLED | WEED GROWTH STAGE | RATE mL/ha | CRITICAL COMMENTS |
|-------------------|------------------------------------|-------------------------------------|-----------------|--|
| 4 to 6 leaf | Cleavers, Fumitory | 2 to 6 leaf | 500 | |
| | Shepherd's purse, Wireweed | | 500 + 5L Asulox | |
| 8 to 10 leaf | Common sowthistle, Prickly lettuce | 2 to 5 leaf | 500 | DO NOT apply Albaugh Fluropix 400 Herbicide to poppies later than the 8 to 10 leaf growth stage as a reduction of alkaloid content could occur. This rate will provide season long control of volunteer potato, but will not control all daughter tubers and will only suppress potatoes over 15 cm tall. |
| | Black nightshade | Cotyledon to 4 leaf | 750 | |
| | Fumitory | 6 to 10 leaf | | |
| | Volunteer potato | From tuber initiation to flower bud | | |

NOT TO USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

TABLE 8: Lucerne (NSW only)

| CROP STAGE GROWTH | WEEDS CONTROLLED | WEED GROWTH STAGE | RATE mL/ha | CRITICAL COMMENTS |
|--|---|-----------------------------|------------|--|
| Established crops at least eighteen months old | Annual ground cherry, Bathurst burr, Noogoora burr, Wild gooseberry | 2 to 8 leaf up to 15cm high | 250 | To minimise crop injury and to maximise weed control, cut, slash or heavily graze the lucerne before application. Wherever possible, irrigate before application to stimulate weed growth. DO NOT treat crops growing on sandy or stony soils. DO NOT treat crops after the summer growing season (after end of March). To broaden the spectrum of weeds controlled, Albaugh Fluropix 400 Herbicide can be mixed with 2,4-DB Amine. |
| | Pigweed | Up to 10 cm diameter | | |

GENERAL INSTRUCTIONS

MINIMUM RECROPPING PERIODS

| PLANT-BACK PERIODS FOR CROPS FOLLOWING THE APPLICATION OF ALBAUGH FLUROPIX 400 HERBICIDE FOR RATES UP TO 750mL/HA | | | |
|---|------|-----|-----|
| RATE ml/ha | 190 | 375 | 750 |
| CROP | DAYS | | |
| Barley, Wheat, Chickpea, Sunflower, Maize, Sorghum | 7 | 7 | 7 |
| Soybean | 7 | 7 | 14 |
| Cotton | 14 | 14 | 28 |

NOTE: Before using Albaugh Fluropix 400 Herbicide in tank mixes with other herbicides, check the plant-back information on all product labels. The time between spraying and planting will be determined by the most residual product, i.e. the product with the longest plant-back period.

MIXING

Albaugh Fluropix 400 Herbicide may be mixed with water or diesel. Mix only sufficient chemical for each day's use and avoid storing.

Mixing in Water: Half fill the spray tank with water and add the required quantity of Albaugh Fluropix 400 herbicide and complete filling. Agitate continuously to ensure thorough mixing before and during application.

Mixing in Diesel: half fill the tank with diesel and add the required quantity of Albaugh Fluropix 400 Herbicide. Add the remainder of the diesel and agitate or shake to mix contents.

Tank mixtures: Wettable powder or dry flowable formulations (e.g. water dispersible granules) should be added to the spray tank first, followed by suspension concentrates (flowables), water soluble salts and then emulsifiable concentrate formulations (Albaugh Fluropix 400 Herbicide). Add spraying oils and surfactants (wettters) last.

OILS AND SURFACTANTS

Oils Use only Uptake Spraying Oil at the rate of 500mL/100L of spray mix. When using less than 100L/ha spray volume, ensure a minimum of 250mL/ha of Uptake is used, unless 1L/100L or 1L/ha is specified.

Surfactants (wettters)

LI to be used at 300mL/100L.

Use a 100% concentrate non-ionic surfactant at 100 mL/100 L of spray mix where required.

COMPATIBILITY

Albaugh Fluropix 400 Herbicide is compatible with the herbicides listed. Follow any regional restrictions, and all directions and restrictions on the label, of any product mixed with Albaugh Fluropix 400 Herbicide.

Atrazine (see below), Glyphosate 360, Metsulfuron methyl (600 g/kg), Glyphosate 450, Broadstrike, cioquintocet-mexyf/clodinafop-propargyl (see below), Eclipse, Tordon 75-D, Diclofop methyl, Tordon 242, Triclopyr (600 g/L), Touchdown, Lontrel, 2,4-D, MCPA, 2,4-DB.

ATRAZINE

AVOID USING HARD WATER WHEREVER POSSIBLE.

Where hard water cannot be avoided, the addition of CALGON water conditioning agent to the spray tank at 100 g/100 L water, before adding any herbicide may improve compatibility.

AGITATION IS VERY IMPORTANT WHEN MIXING Albaugh Fluropix 400 Herbicide and ATRAZINE.

Albaugh Fluropix 400 Herbicide plus atrazine tank mixes must be agitated vigorously and continuously during mixing and application. After mixing DO NOT allow to stand without agitation. Ensure that the time from mixing to the end of application is not more than 2 hours. If settling out occurs re-suspension is difficult, even with vigorous agitation. Agitation using only the pump's by-pass is usually inadequate, particularly with larger tanks (more than 2000 L). Additional mechanical agitation will be necessary in large tanks, computer sprayers and mixing tanks. When additional surfactant is required, add a 100% concentrate non-ionic surfactant at 100 mL/100 L of spray mix. DO NOT use a spraying oil when tank mixing Albaugh Fluropix 400 Herbicide and atrazine.

| GUIDELINES FOR TANK-MIXING ALBAUGH FLUROPIX 400 HERBICIDE AND ATRAZINE FORMULATIONS | | | | | | |
|---|--------------|----------------|------------------|-----------------------------|------------|---|
| TANK MIX | RATE (/Ha) | WATER HARDNESS | | MINIMUM WATER VOLUME (L/HA) | | CRITICAL COMMENTS |
| | | Soft | Hard(1026ppm ca) | Ground | Aerial | |
| Albaugh Fluropix 400 Herbicide + Atrazine (900g/kg) WG Herbicide | 375mL +1.1kg | √ | √ | 100 | Do not use | Sediment may be difficult to resuspend and may block nozzels. |
| Albaugh Fluropix 400 Herbicide +Atrazine (600g/L) | 375mL + 1.7L | √ | √ | 100 | Do not use | |

Cloquintocet-Mexyl/Clodinafop-Propargyl

Always use Uptake Spraying Oil with Albaugh Fluropix 400 Herbicide + Cloquintocet-mexyl/clodinafop-propargyl tank-mixes at 500 mL/100 L of spray mix with a minimum of 250 mL/ha. DO NOT mix Albaugh Fluropix 400 Herbicide with Cloquintocet-mexyl/clodinafop-propargyl if the grass weeds are not actively growing. Always use the maximum label rate of Cloquintocet-mexyl/clodinafop-propargyl for the appropriate grass growth stage. DO NOT use Albaugh Fluropix 400 Herbicide at more than 0.75 L/ha in tank mixes with Cloquintocet-mexyl/clodinafop-propargyl.

GLYPHOSATE 450

When mixing Albaugh Fluropix 400 Herbicide with Glyphosate 450 to control both grass and broadleaf weeds, refer to the Glyphosate 450 label for use rates and adjuvants recommended for the grasses. DO NOT use Glyphosate 450 at less than 1.2 L/ha in tank mixes with Albaugh Fluropix 400 Herbicide, when barnyard grass, buttongrass, crowsfoot grass, native millet and liverseed grass are the target species.

APPLICATION METHODS and WATER RATES

BROADCAST APPLICATION IN CROPPING, PASTURE AND FALLOW SITUATIONS A. Ground application (Boom) Apply Albaugh Fluropix 400 Herbicide with an accurately calibrated boom sprayer, in at least 50 L/ha water (100-400 L/ha for sugar cane). Flat fan nozzles are recommended using pressures in the range 200 to 300kPa.

Set the boom at a height to ensure a double overlap of the nozzle patterns.

B. Ground directed application (Dropper nozzles)

To minimise crop effects, dropper nozzles should be used in sorghum when the crop is beyond the 8 leaf growth stage and in maize and sweet corn when the crop is beyond the 6 leaf growth stage. Adjust the nozzles to direct the spray into the base of the crop and away from the leaves and the growing point. See manufacturer's directions for setting up and calibration of dropper nozzles

C. Aerial application

Apply in a minimum volume of at least 35 L/ha water (60 L/ha in sugarcane).

Use equipment calibrated to produce droplets with an average diameter (Volume Mean Diameter; VMD) of 250 – 350 microns. DO NOT apply when the temperature is above 30^o C, when there is no wind or when the wind is blowing toward susceptible crops. DO NOT use human flaggers unless they are protected by engineering controls such as enclosed cabs.

WOODY WEED SITUATIONS

Weeds must be actively growing to attain optimal effect. Delay the treatment of regrowth following bulldozing, slashing, burning, ploughing or a previous chemical treatment until it has at least 1 metre of new, vigorous, growth.

A. High Volume Application

Hand Gun

Apply the recommended mix to obtain full coverage of leaves and stems using a number 6 – 8 tip at 700 to 1500 kPa. To obtain good coverage, a spray volume of 1500 to 4000 L/ha (15 to 40 L/100m²) is required per infested hectare. Ensure thorough coverage to the point of runoff.

Knapsack

Knapsack sprayers may be used on smaller infestations where penetration and coverage of the canopy is easier to achieve. Use the same use rate and spray techniques as for handgun application.

B. Low Volume, High Concentrate Application

Drench Gun or Gas-Powered Gun

Apply the recommended mixture uniformly across the foliage by applying 50mL shots to cover 4 to 5 m² of surface area of plant. This is approximately equivalent to 20 droplets per cm² of the leaf surface. Use a marking agent as recommended by the equivalent manufacturer to check spray coverage.

C. Basal Bark and Cut Stump Application

Basal Bark

DO NOT apply to wet stems as this can repel the diesel mixture. Spray or paint the recommended mixture around the base of each stem from ground level to a height of at least 30 cm from the ground, wetting the bark to the point of runoff. Apply with a paint brush or a pressure sprayer with an appropriate lance and solid cone nozzle. If using spray equipment use low pressures (\leq 200 kPa) sufficient to form a cone of spray.

Old rough bark will require more spray than smooth or young thin bark.

Cut Stump

Apply the recommended mixture liberally to the freshly cut stump immediately after cutting. Apply by spraying or painting the cut surface and sides of the stump.

Best results are obtained when the stems are cut less than 15 cm above the ground.

CLEANING SPRAY EQUIPMENT

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto wasteland away from desirable plants and water courses.

Cleaning equipment after using water-based sprays:

Rinsing: After using Albaugh Fluropix 400 Herbicide, empty the tank completely and drain the whole system.

Thoroughly wash inside the spray unit using a pressure hose. Drain and clean any filters in the tank, pump, lines, hoses and nozzles.

After cleaning the tank as above, quarter fill the clean water and circulate through the pump, lines and nozzles. Drain and repeat the rinsing procedure twice.

Decontamination (before spraying cotton and other sensitive crops; see PROTECTION OF CROPS): Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. Liquid SURF, OMO, DRIVE) at 500 mL/100L of water or the powder equivalent at 500 g/100 L and circulate throughout the system for at least fifteen minutes.

Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

Cleaning equipment after using diesel – based sprays:

On completion of spraying, use a degreaser such as Caltex Kwik-D-Grease to remove traces of diesel from the sprayer. Rinse tank and spray through nozzles with water to remove degreaser. Then quarter fill the tank and add an alkali detergent (e.g. liquid SURF, OMO, DRIVE) at 50 mL/10L of water or the powder equivalent at 50 g/10 L.

Shake sprayer to circulate the washing solution throughout the sprayer, then spray the solution through the nozzles. Rinse well with clean water to remove the detergent. To clean brushes and containers, spray liberally with degreaser. Hose off with clean water and repeat using detergents as above. DO NOT use this equipment for any other purpose.

SMALL SPILL MANAGEMENT

Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb the wash liquid for disposal.