



Product Name: Albaugh Ditto Selective Herbicide
APVMA Approval No.: 81130/141541

Label Name:	Albaugh Ditto Selective Herbicide
--------------------	-----------------------------------

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
-------------------------	---

Constituent Statements:	ACTIVE CONSTITUENTS: 240 g/L CLODINAFOP-PROPARGYL 60 g/L CLOQUINTOCET-MEXYL SOLVENTS: 575 g/L LIQUID HYDROCARBONS 100 g/L N-METHYL-2-PYRROLIDONE
--------------------------------	---

Mode of Action:	GROUP 1 HERBICIDE
------------------------	--------------------------

Statement of Claims:	For the control of Wild Oats, Paradoxa Grass (Annual Phalaris), Canary Grass and Annual Ryegrass in Wheat. Highly active on Wild Oats and can be mixed with many broadleaf herbicides.
-----------------------------	--

Net Contents:	1L, 5L
----------------------	--------

Restraints:	DO NOT apply if rainfall is expected within 2 hours. DO NOT apply to weeds or crops which are under stress. For example, due to very dry, very wet, frosty or diseased conditions, nutrient deficiency or high insect pressure. DO NOT apply to weeds or crops immediately after a prolonged period of frosts. DO NOT apply under poor growing conditions.
--------------------	---

Directions for Use:	This section contains file attachment.
----------------------------	--

Other Limitations:	
---------------------------	--

Withholding Periods:	<p>HARVESTED GRAIN: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.</p> <p>ANIMAL GRAZING/FEEDING: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.</p>
-----------------------------	---

Trade Advice:	
----------------------	--

General Instructions:	<p>Albaugh Ditto Selective Herbicide should be applied only when weeds are actively growing.</p> <p>MIXING Fill the spray tank to one quarter full. Then add Albaugh Ditto Selective Herbicide while adding water to make up the final spray volume. Pour Albaugh Ditto Selective Herbicide into the stream of incoming water. Good agitation at the time Albaugh Ditto Selective Herbicide is added to the tank will ensure good mixing. Add the required amount of D-C-Trate*, Hasten* or Uptake* oil just before the tank is full of water and with the agitators in motion.</p> <p>Mixing order Some products may react with other products if they are not mixed in the correct order. The general mixing order of products should be –</p> <ol style="list-style-type: none"> 1. Water conditioners or buffers 2. Water dispersable granules (WG) 3. Wettable powders (WP) 4. Flowable or suspension concentrates (SC) 5. Emulsifiable concentrates (EC) (Albaugh Ditto Selective Herbicide) 6. Water based or soluble concentrates 7. Oils and wetters - eg Hasten, BS 1000 <p>It is important to ensure that each individual component of the tank mix is fully dissolved and in solution before the next product is added to the tank mix, otherwise mixing problems may occur.</p> <p>Minimum Oil Volumes The minimum amount of oil to be applied with Albaugh Ditto Selective Herbicide will depend on the type of oil used. When applying Albaugh Ditto Selective Herbicide using low water volumes, DO NOT use less than the minimum oil volumes specified below:</p> <ul style="list-style-type: none"> • For oils normally applied at 1% v/v (eg D-C-Trate), DO NOT apply less than 500 mL oil/ha. • For oils normally applied at 0.5% v/v (eg Uptake, Hasten), DO NOT apply less than 250 mL oil/ha. <p>CLEANING SPRAY EQUIPMENT After using Albaugh Ditto Selective Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.</p> <p>To rinse: After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.</p> <p>To decontaminate: Before spraying cereals (except wheat), maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Drain and repeat the rinsing procedure. Then quarter fill the tank and add a liquid alkali detergent (eg SURF*, OMO*, DRIVE*) at 0.5 L/100 L of water and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove the filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain. Dispose of all water used for cleaning in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots.</p> <p>APPLICATION Ground Application: Boom spray with flat fan nozzles. Ensure good spray coverage is</p>
------------------------------	--

	<p>obtained. Apply using 50 to 110 L water per hectare. The ideal droplet size is 200 to 300 microns VMD.</p> <p>Aircraft Application: For best results use 20 to 30 L water per hectare and spray at 2m to 3m above the crop. Ideal droplet size is 250 to 350 microns VMD. For rotary atomizers (micronairs) this can be achieved using blade angles of 55° to 65°. Where booms and nozzles are attached, flat fan nozzles with a spray angle of 65° to 90° should be used. Avoid applying Albaugh Ditto Selective Herbicide if wind speeds are greater than 5m/s. If spraying in calm conditions use larger droplets and reduce flying height.</p>
--	---

Resistance Warning:	<p>Albaugh Ditto Selective Herbicide is a member of the aryloxyphenoxypropionate ("fop") group of herbicides. The product has the inhibition of fat (lipid) synthesis, (or inhibitors of acetyl CoA carboxylase) mode of action. For weed resistance management, the product is a Group 1 herbicide.</p> <p>Some naturally occurring weed biotypes resistant to the product and other Group 1 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 1 herbicides.</p> <p>Since the occurrence of resistant weeds is difficult to detect prior to use, Albaugh Asia Pacific Limited accepts no liability for any losses that may result from the failure of this product to control resistant weeds.</p>
----------------------------	--

Precautions:	<p>Re-entry Period DO NOT enter treated areas without protective clothing until the spray has dried.</p>
---------------------	--

Protections:	<p>PROTECTION OF CROPS, NATIVE AND OTHER NON - TARGET PLANTS DO NOT apply on or near shrubs, trees, lawns or crops other than wheat. DO NOT drain or flush equipment on or near desirable trees or other plants, where their roots may extend, or in situations where by movement of soil or by seepage absorption of the herbicide may occur. DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. DO NOT allow spray to drift onto adjacent fallow land.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT apply to irrigation drains or channels. DO NOT contaminate dams, streams, creeks and rivers with the chemical, spray mixture, used containers or equipment washings.</p>
---------------------	--

Storage and Disposal:	<p>STORAGE AND DISPOSAL Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.</p> <p>Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
------------------------------	--

Safety Directions:	<p>Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid inhaling spray mist. Avoid contact with eyes and skin. When opening the</p>
---------------------------	---

container, preparing spray and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat, elbow-length butyl rubber gloves and face shield or goggles. If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**First Aid
Instructions:**

If poisoning occurs contact a doctor or Poisons Information Centre.
Phone Australia 131 126.
If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

First Aid Warnings:

DIRECTIONS FOR USE

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE	WEED GROWTH STAGE	CRITICAL COMMENTS	
Wheat 2 leaf to late jointing (Zadocks 12-37)	Wild Oats	Qld, Nthn NSW only	65 mL + 1 L of D-C- Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 to 5 leaf	NOT TO BE MIXED WITH BROADLEAF HERBICIDES AT THESE RATES Apply at 85 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.	
			85 mL + 1 L of D-C- Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 leaf to mid tillering		
			85 mL + 1 L of D-C- Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 to 5 leaf		
			125 mL + 1 L of D-C- Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 leaf to mid tillering		
			Sthn NSW, Vic, SA, WA only	65 mL + 1 L of D-C- Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water.	2 to 5 leaf	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing Albaugh Ditto Selective Herbicide with certain broadleaf weed herbicides some reduction in grass weed control may occur. Refer to compatibility chart for specific rates for tank mixes. Unless otherwise specified always use D-C Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 85 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
				85 mL + 1 L of D-C- Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water.	2 leaf to mid tillering	
	Paradoxa Grass (<i>Phalaris paradoxa</i>) Canary Grass (<i>Phalaris minor</i>)	Qld, NSW, Vic, SA only	85 mL + 1 L of D-C Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 to 3 leaf	NOT TO BE MIXED WITH BROADLEAF HERBICIDES AT BELOW 160 mL/ha Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing Albaugh Ditto Selective Herbicide with certain broadleaf weed herbicides some reduction in grass weed control may occur. Apply at 160 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.	
			125 mL + 1 L of D-C- Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 to 5 leaf		
160 mL + 1 L of D-C- Trate oil or 500 mL Hasten or Uptake per 100 L of water.			2 leaf to early tillering			
Annual Ryegrass	NSW, Vic, SA, WA only	160 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 to 4 leaf	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing Albaugh Ditto Selective Herbicide with certain broadleaf weed herbicides some reduction in grass weed control may occur. Apply at 210 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.		
		210 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 leaf to early tillering			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.