



**Company Name:** SHANDONG RAINBOW INTERNATIONAL CO., LTD.  
**Product Name:** I-MOX 700 WG HERBICIDE  
**APVMA Approval No:** 81174/102487

<b>Label Name:</b>	I-MOX 700 WG HERBICIDE
--------------------	------------------------

<b>Signal Headings:</b>	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
-------------------------	--

<b>Constituent Statements:</b>	ACTIVE CONSTITUENT: 700 g/kg IMAZAMOX
--------------------------------	---------------------------------------

<b>Mode of Action:</b>	<table border="1"><tr><td>GROUP</td><td><b>B</b></td><td>HERBICIDE</td></tr></table>	GROUP	<b>B</b>	HERBICIDE
GROUP	<b>B</b>	HERBICIDE		

<b>Statement of Claims:</b>	For the post-emergence control of certain annual grass and broad leaved weeds in field peas, legume-based pastures, lucerne, peanuts and soybeans, as specified in the Directions for Use table.
-----------------------------	--

<b>Net Contents:</b>	1kg 2kg 2.5kg 3kg 5kg 10kg 15kg 20kg
----------------------	---

<b>Restrains:</b>	DO NOT apply to crops or weeds stressed by factors such as root or foliar diseases, water logging, nutrient deficiencies or extremes of temperature and moisture. DO NOT apply to crops stressed by any previous herbicide treatments.
-------------------	---

<b>Directions for Use:</b>	This section contains file attachment. <b>File Name:</b> DIRECTIONS FOR USE .pdf <b>File Size:</b> 77323 bytes
----------------------------	--

<b>Other Limitations:</b>	
---------------------------	--

<b>Withholding Periods:</b>	<p>GRAZING          Lucerne, Pastures: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.          Peanuts, Soybeans: DON NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.          Field Peas: DO NOT GRAZE OF CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.          HARVEST          Field peas, Peanuts, Soybeans: NOT REQUIRED WHEN USED AS DIRECTED.</p>
-----------------------------	--

<b>Trade Advice:</b>	
----------------------	--

<b>General Instructions:</b>	<p>This section contains file attachment.  <b>File Name:</b> GENERAL INSTRUCTIONS.pdf  <b>File Size:</b> 38481 bytes</p>
------------------------------	--

<b>Resistance Warning:</b>	<p>RESISTANT WEEDS WARNING          GROUP B HERBICIDE          I-Mox 700 WG Herbicide is a member of the imidazoline group of herbicides. This product has the ALS mode of action. For weed resistance management I-Mox 700 WG Herbicide is a Group B Herbicide. Some naturally occurring weed biotypes resistant to the product and other Group B Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group B Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Shandong Rainbow International Co., Ltd accepts no liability for any losses that may result from failure of this product to control resistant weeds.</p>
----------------------------	--

<b>Precautions:</b>	<p>This section contains file attachment.  <b>File Name:</b> PRECAUTIONS.pdf  <b>File Size:</b> 52909 bytes</p>
---------------------	---

<b>Protections:</b>	<p>This section contains file attachment.  <b>File Name:</b> PROTECTION STATEMENTS.pdf  <b>File Size:</b> 44100 bytes</p>
---------------------	---

<b>Storage and Disposal:</b>	<p>STORAGE AND DISPOSAL          Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.</p>
------------------------------	--

<b>Safety Directions:</b>	<b>SAFETY DIRECTIONS</b> Will irritate the eyes. When opening the container and preparing spray, wear face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash face shield or goggles.
---------------------------	---

<b>First Aid Instructions:</b>	<b>FIRST AID</b> If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
--------------------------------	--

<b>First Aid Warnings:</b>	
----------------------------	--

**DIRECTIONS FOR USE  
RESTRAINTS**

Do NOT apply to crops or weeds stressed by factors such as root or foliar diseases, water logging, nutrient deficiencies or extremes of temperature and moisture.

Do NOT apply to crops stressed by any previous herbicide treatments.

CROP	WEEDS CONTROLLED	RATE/Ha	WHP	CRITICAL COMMENTS
Field Peas Post-emergence See "Crop Safety" re varieties	Barley grass ( <i>Hordeum leporinum</i> ), Brome grass ( <i>Bromus</i> spp.), Deadnettle ( <i>Lamium amplexicaule</i> ), Indian Hedge mustard ( <i>Sisymbrium orientale</i> ), Storksbill ( <i>Erodium</i> spp.), Turnip weed ( <i>Rapistrum rugosum</i> ), Volunteer barley ( <i>Hordeum vulgare</i> ), Volunteer oats ( <i>Avena sativa</i> ), Volunteer lupins ( <i>Lupinus</i> spp.), Volunteer triticale ( <i>Triticosecale</i> spp.), Volunteer wheat ( <i>Triticum aestivum</i> ), Wild Oats ( <i>Avena</i> spp.), Wild Turnip ( <i>Brassica tournefortii</i> ), Doublegee ( <i>Emex australis</i> ), *Shepherds purse ( <i>Capsella bursa-pastoris</i> ), * Threehorn bedstraw ( <i>Calium tricomutum</i> ), *Wild radish ( <i>Raphanus raphanistrum</i> ), *Wireweed ( <i>Polygonum aviculare</i> )	45g plus BS 1000, or equivalent, at 200mL per 100L water	6 weeks (grazing)	<p><b>DO NOT apply to field peas beyond the 4-node stage.</b> DO NOT use adjuvants other than BS1000 or equivalent. Apply to actively growing broadleaved weeds in the cotyledon to 3-leaf stage. Apply to grass weeds up to the 2-tiller stage. Good crop growth with aid weed control. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. *Control will not be 100% but surviving plants will generally be retarded and will not complete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds. Leaf yellowing and height suppression may occur, especially under adverse growing conditions such as moisture stress. Flowering may be delayed.</p> <p><b>IMPORTANT: Refer to CROP SAFETY section of label re use on field peas.</b></p> <p>Refer to FOLLOW CROPS section of this label regarding follow crops.</p>
Legume-based pastures Post-emergence Clovers, Lucerne, Medics, Serradellas only; see "Crop Safety" re varieties		45g plus Hasten™ or Kwicken™ at 500mL per 100L water	7 days (grazing)	<p><b>Seedling</b> - Apply to legumes after full emergence of the third trifoliolate leaf but well before commencement of flowering. <b>Established</b>- Apply before commencement of flowering. Apply to actively growing broadleaved weeds in the cotyledon to 3-leaf stage. Apply to grass weeds up to the 2-tiller stge. Good pasture growth will aid weed control. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Where pasture stand is poor, weed control may be inadequate due to lack of competition. Ensure sufficient spray coverage of weeds particularly in dense pastures. *Control will not be 100% but surviving plants will generally be retarded and will not complete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds. Refer to CROP SAFETY section of this label re varieties of clovers and medics and other pasture species. Note: use of the 50g/ha rate plus Hasten or Kwicken may severely damage medics. Refer to FOLLOW CROPS section of this label regarding follow crops.</p>
50g plus Hasten or Kwicken, at 500mL per 100L water plus Spraymater Liase at 2L per 100L water				

CROP	WEEDS CONTROLLED	RATE/Ha	WHP	CRITICAL COMMENTS
Lucerne Post-emergence (See "Crop Safety" re varieties)	As for post emergence use in field peas (winter weeds) and soybeans (summer weeds)	45 or 50g plus BS 1000, or equivalent, at 200mL per 100L water plus BOOST at 2L per 100L water	7 days (grazing)	Apply to actively growing weeds at growth stages as for field peas (winter weeds) and soybeans (summer weeds).  DO NOT use the Hasten or Kwicken mix on seedling lucerne. Use the 50g rate for summer weeds.  Crop Stage: Seedling: Apply when crop is at 2 trifoliate leaf stage and before commencement of flowering. Established: Apply as above following cutting or grazing.  Refer to CROP SAFETY section of this label re varieties of lucerne
Post-emergence Peanuts, Soybeans	Amaranth ( <i>Amaranthus</i> spp.), Barley grass ( <i>Hordeum leporinum</i> ), Barnyard grass ( <i>Echinochloa crus-gali</i> ), Bellvine ( <i>Ipomoea plebeia</i> ), Brome grass ( <i>Bromus</i> spp.), Deadnettle ( <i>Lamium amplexicaule</i> ), Fathen ( <i>Chenopodium album</i> ), Fierce Thornapple ( <i>Datura ferox</i> ). Indian Hedge mustard ( <i>Sisymbrium orientale</i> ), Liverseed grass ( <i>Urochloa panicoides</i> ), Turnip weed ( <i>Rapistrum rugosum</i> ), Volunteer barley ( <i>Hordeum vulgare</i> ), Volunteer oats ( <i>Avena sativa</i> ), Volunteer lupins ( <i>Lupinus</i> spp.), Volunteer triticale ( <i>Triticosecale</i> spp.), Volunteer wheat ( <i>Triticum aestivum</i> ), Wild gooseberry ( <i>Physalis minima</i> ), Wild Oats ( <i>Avena</i> spp.), Wild Turnip ( <i>Brassica tournefortii</i> ), *Anoda weed ( <i>Anoda cristata</i> ), *Awnless barnyard grass ( <i>Echinochloa colona</i> ), *Blackberry nightshade ( <i>Solanum nigrum</i> ), *Caltrop ( <i>Tribulus terrestris</i> ), *Chickweed ( <i>Stellaria media</i> ), *Crabgrass ( <i>Digitaria ciliaris</i> ), *Doublegee ( <i>Emex australis</i> ), *Noogoora burr ( <i>Xanthium pungens</i> ), *Shepherds purse ( <i>Capsella bursa-pastoris</i> ), * Threehorn bedstraw ( <i>Calium tricomutum</i> ), *Wild radish ( <i>Raphanus raphanistrum</i> ), *Wireweed ( <i>Polygonum aviculare</i> )	50g plus <b>EITHER</b> BS 1000 (or equivalent), at 200mL/100L water plus BOOST at 2L/100L water <b>OR</b> Hasten or Kwicken, at 500mL or 1L /100L water	4 weeks (grazing)	DO NOT apply to crops beyond the 6-leaf stage. Apply to actively growing weeds in the cotyledon to 2-leaf stage. Apply to grass weeds up to the 2-tiller stage.  If using the Hasten or Kwicken mix, use the 1L/100L rate where grass weeds predominate.  *Control will not be 100% but surviving plants will generally be retarded and will not compete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds.  Refer to FOLLOW CROPS section this label regarding follow crops.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## **GENERAL INSTRUCTIONS**

This product is for early post-emergence weed control in field peas, lucerne, legume-based pastures, soybeans and peanuts. Weeds will either die or remain stunted and will not compete with the crop. For weeds marked with an asterisk in the above table, the level of control will vary significantly depending on factors such as climatic conditions following application, crop vigour, weed seed depth, etc.

A spray adjuvant must always be added, as per the "Directions for Use" table, while Boost liquid ammonium sulphate should also be added in situations indicated in the table.

Good crop competition is essential for effective weed control.

## **MIXING**

I-Mox 700 WG Herbicide is a water dispersible granular formulation. Part fill the spray tank with water, then with the agitator running, add the required amount of product, then fill the tank with water. When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product.

## **APPLICATION**

For ground application only. Apply with boom equipment in not less than 50L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands.

DO NOT apply by aircraft.

DO NOT apply by mister.

I-Mox 700 WG Herbicide should be applied a minimum of two hours before rainfall or irrigation.

DO NOT apply I-Mox 700 WG Herbicide more than once per growing season.

## **EQUIPMENT CLEANUP**

Thoroughly flush all spray equipment with water following the use of I-Mox 700 WG Herbicide and before use with other products. Rinse water should NOT be discharged where it will reach streams, water bodies or natural vegetation.

## **COMPATIBILITY**

This product is compatible with MCPA, Simazine, Dimethoate, alpha-cypermethrin, phosmet and omethoate.

DO NOT tank mix with selective post-emergence grass herbicides. DO NOT apply these herbicides following use of I-Mox 700 WG Herbicide until grasses have resumed active growth. Alternatively, apply the grass herbicide at least one day before using I-Mox 700 WG Herbicide.

DO NOT tank mix with either MCPA Amine or 2,4-D Amine when using in pastures.

## FOLLOW CROPS

Under conditions, such as very dry season, which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible. The following minimum re-cropping intervals (months after application) should be observed.

### Following use in lucerne, legume-based pastures and field peas:

Months after application		
0	10	21
Field peas Canola varieties with CLEARFIELD technology Maize varieties with CLEARFIELD technology Wheat varieties with CLEARFIELD technology	Chickpeas Faba Beans Lucerne Lupins Pasture Legumes Vetch *Barley *Wheat (except varieties with CLEARFIELD technology) *Triticale	All other crops including canola (except varieties with CLEARFIELD technology) Oats Safflower

\*The following additional requirements apply if it is intended to sow WHEAT, BARLEY or TRITICALE during the next winter season:

DO NOT apply I-Mox 700 WG Herbicide later than the end of August.

DO NOT use I-Mox 700 WG Herbicide in areas where rainfall from time of spraying to sowing of cereals is expected to be below 200mm.

Furthermore:

DO NOT use on soils of pH 5.5 (CaCl<sub>2</sub>) or less in areas where rainfall from spraying to sowing of cereals is expected to be below 300mm.

In late, short seasons where the soil is cold for most of the time it is wet, break-down will be slower and plant-back times will be extended.

If expected rainfall is not received following use of I-Mox 700 WG Herbicide, consult your local Shandong Rainbow representative before planting wheat, barley or triticale. (In calculating rainfall actually received, exclude single, isolated falls which do not result in periods of continuous soil moisture to allow microbial breakdown to occur in the root zone).

### Following use in summer crops.

Irrigated only

Months after application			
0	5	10	18
Mung beans Peanuts Soybeans Canola varieties with CLEARFIELD technology Maize varieties with CLEARFIELD technology Wheat varieties with CLEARFIELD technology	*Chickpeas *Lucerne *Lupins *Pasture Legumes *Barley *Wheat (except varieties with CLEARFIELD technology)	**Maize (except varieties with CLEARFIELD technology) **Sorghum **Cotton **Oats **Sunflower	All other crops (providing rainfall and irrigation exceeds 2000mm)

\*DO NOT plant these crops unless interim rainfall (rainfall plus irrigation) from application to sowing is at least 500mm.

\*\* DO NOT plant these crops unless interim rainfall (rainfall plus irrigation) from application to sowing is at least 800mm.

Dryland only

Months after application				
0	8	10	15	22
Mung beans Peanuts Soybeans Canola varieties with CLEARFIELD technology Maize varieties with CLEARFIELD technology Wheat varieties with CLEARFIELD technology	*Lucerne *Barley *Wheat (except varieties with CLEARFIELD technology)	**Chickpeas **Maize (except varieties with CLEARFIELD technology) **Sorghum **Cotton **Oats **Sunflower	Pasture Legumes Lupins	All other crops (providing rainfall and irrigation exceeds 2000mm)

\*DO NOT plant these crops unless interim rainfall (rainfall plus irrigation) from application to sowing is at least 500mm.

\*\*DO NOT plant these crops unless interim rainfall (rainfall plus irrigation) from application to sowing is at least 800mm.



## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT spray within 50m of wetlands or waterways.

### **Crop Safety:**

This product may cause shortening of plant internodes and may, in some circumstances, lead to transient crop yellowing but plants will soon recover. This effect is more pronounced under poor growth conditions including conditions of prolonged moisture stress.

### **Field Peas**

DO NOT apply to field peas beyond the 4 node stage.

DO NOT apply if frost is forecast.

DO NOT use I-Mox 700 WG Herbicide after simazine (unregistered in field peas) and DO NOT exceed 250g ac/ha diuron if use of I-Mox 700 WG Herbicide is planned.

DO NOT roll field peas after germination if use of I-Mox 700 WG Herbicide is planned.

Before using this product on field pea varieties other than Bohatyr, Bonzer, Dun, Dundale, Glenroy, Laura, Magnet, Mukta, Santi and Wirrega, contact Shandong Rainbow International Co., Ltd for up-to-date information. DO NOT use on Alma, Excell and Parafield varieties.

### **Pasture Legumes and Perennial Grasses**

Safety to medic varieties may be lower than to clover varieties. Safety to seedling pastures less than once year old may be lower than to older established pastures. Transient biomass reduction and leaf yellowing may occur to pasture legumes and perennial grass, with good recovery occurring under favourable seasonal conditions. In some situations a reduction in perennial grass numbers may occur as a result of unfavourable seasonal conditions following application. For up-to-date information on all pasture clover and medic varieties and other temperate pasture species, contact Shandong Rainbow International Co., Ltd before use.

### **Lucerne**

Before using this product on lucerne varieties other than Hunterfield, Sceptre, Sequel, Siriver and Trifecta, contact Shandong Rainbow International Co., Ltd for up-to-date information.

**DO NOT USE THE HASTEN OR KWICKIN MIX ON SEEDLING LUCERNE.**

### **Replanting**

If replanting is necessary in a field previously treated with I-Mox 700 WG Herbicide, the field may be replanted to field peas (above listed varieties only), soybeans or peanuts. DO NOT apply a second treatment of I-Mox 700 WG Herbicide.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Toxic to aquatic flora. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.