



Product Name: STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT
APVMA Approval No: 81483/121704

Label Name:	STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT
-------------	--

Signal Headings:	FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	50 g/L DICYCLANIL
-------------------------	-------------------

Claims:	For the protection of sheep, either off-shears or with any length wool, for up to 18 to 24 weeks against strike by dicyclanil-susceptible strains of blowflies (<i>Lucilia cuprina</i>) when applied by spraying. For the protection of mulesing and marking wounds on sheep against fly strike (<i>Lucilia cuprina</i>) during the wound healing process. READ DIRECTIONS FOR USE BEFORE OPENING OR USING THIS PRODUCT
---------	--

Net Contents:	10L 15L 1L 20L 22L 5L
---------------	--------------------------------------

Directions for Use:	
---------------------	--

Restrains:	DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption. Re-treatment Interval: DO NOT re-treat animals for a minimum of 35 days after last treatment.
------------	--

Contraindications:	
--------------------	--

Precautions:	
--------------	--

Side Effects:	
Dosage and Administration:	<p>The dose volume is determined by the weight of sheep being treated. Apply STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT according to the diagrams and tables below. DO NOT DILUTE. Apply directly from the container or an appropriate back pack using a spray-on applicator gun recommended by Jurox.</p> <p>STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT is applied as a low-volume treatment onto areas where protection is required. When applying STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT, the nozzle of the applicator gun should be high enough above the fleece (approximately 15 – 25 cm), to achieve a 15 cm wide band at each pass for protection of body strike. For treatment of mulesing wounds, the band must be wide enough to cover the wound and 25 mm onto the surrounding wool.</p> <p>This section contains file attachment. File Name: Dose table for e-label.pdf File Size: 46750 bytes</p>
General Directions:	<p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p> <p>STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT provides season long protection to sheep against body, breech and poll strike and may be used either off-shears or on sheep with any length wool.</p> <p>STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT also protects mulesing and marking wounds on lambs against fly strike during the wound healing process.</p> <p>STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT may be applied to all breeds of sheep to obtain protection against fly strike.</p> <p>DO NOT apply during gusty or windy weather as spray drift may reduce the period of protection. It is good agricultural practice for mulesing / marking to be carried out at times of minimal fly activity. Reduced efficacy may occur in sheep affected with dermatophilosis. Significant soiling with dags, mud, dust or urine may reduce efficacy. Where significant dags are present, a light crutching is indicated prior to treatment. Protection period may be reduced if sheep scour heavily following application.</p>
Withholding Periods:	<p>MEAT: DO NOT USE less than 14 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.</p> <p>WOOL HARVESTING INTERVAL: DO NOT USE less than 3 months before shearing or fibre collection.</p> <p>Use of this product may result in wool residues that may not comply with European Union environmental quality standards.</p> <p>Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.</p>
Trade Advice:	<p>EXPORT SLAUGHTER INTERVALS (ESIs):</p> <p>SHEEP - DO NOT USE less than 35 days before slaughter for export.</p> <p>Before using this product, confirm the current ESIs from Jurox Pty Limited on 1800 023 312 or the APVMA website (www.apvma.gov.au/residues).</p>

Safety Directions:	Harmful if swallowed. May irritate the skin. Will damage eyes. Avoid contact with eyes and skin. When opening the container and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length rubber gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and goggles and contaminated clothing.
--------------------	--

First Aid Instructions:	First Aid is not generally required. If in doubt, contact a Poisons Information Centre (Phone Australia 13 11 26) or a doctor.
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	<p>ADDITIONAL USER SAFETY INFORMATION</p> <p>Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling.</p> <p>DO NOT USE less than 3 months before shearing or fibre collection</p> <p>SAFETY DATA SHEET: Additional information is listed in the Safety Data Sheet which is available on the Jurox website (www.jurox.com.au) or by contacting 1800 02 3312</p>
-------------------------	--

Environmental Statements:	<p>ENVIRONMENTAL PROTECTION</p> <p>DO NOT contaminate wetlands or watercourses with this product or used containers.</p>
---------------------------	---

Disposal:	<p>[Pack sizes less than or equal to 1 L] Dispose of container by wrapping in paper and putting in garbage.</p> <p>[Pack sizes greater than 1 L] Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
-----------	--

Storage:	KEEP OUT OF REACH OF CHILDREN. Store below 30 °C (room temperature). Protect from light. Store in original container, tightly closed in a cool, dry place. DO NOT re-use the container.
----------	---

Dose Table for STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT

FOR THE PREVENTION OF BODY STRIKE (<i>Diagram 1</i>)					CRITICAL COMMENTS
SHEEP WEIGHT (kg)	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	NO. OF SHEEP TREATED PER [1 L][5 L] [10 L][15 L][20 L][22 L]	1. Use only a spray-on applicator gun recommended by Jurox. 2. Ensure the applicator gun is clean. 3. Calibrate applicator gun using an accurate volume measure. 4. Ensure all areas to be protected are treated. 5. When preventing body strike ensure bands overlap along the midline – see Diagram 1.
10 - 20	14	7	2	[71] [357] [714] [1071] [1428] [1571]	
21 - 30	16	8	2	[62] [312] [625] [937] [1250] [1375]	
31 - 50	20	10	2	[50] [250] [500] [750] [1000] [1100]	
> 50	24	12	2	[41][208][416] [625][833][916]	
FOR THE PREVENTION OF BREECH STRIKE (<i>Diagram 1</i>)					<i>Diagram 1 (Positions of bands on sheep for prevention of body, breech and poll strike)</i>
SHEEP WEIGHT (kg)	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	NO. OF SHEEP TREATED PER [1 L][5 L] [10 L][15 L][20 L][22 L]	6. When protecting mulesing wound from fly strike, take special care to treat the wound and the surrounding 25 mm of wool – see Diagram 2.
10 - 20	7	7	1	[142][714][1428] [2142][2857][3142]	
21 - 30	8	8	1	[125][625][1250] [1875][2500][2750]	
31 - 50	10	10	1	[100][500][1000] [1500][2000][2200]	
> 50	12	12	1	[83][416][833] [1250][1666][1833]	<i>Diagram 2 (Area of sheep to treat after mulesing)</i>
FOR THE PREVENTION OF POLL STRIKE (<i>Diagram 1</i>)					7. When protecting marking wounds from fly strike, take special care to treat the wound and surrounding wool – see diagram 3.
Rams and sheep susceptible to poll strike should be treated. Apply 5 mL of STRIKEFORCE-S SPRAY-ON SHEEP BLOWFLY TREATMENT to the areas requiring protection, around the poll and horns.					
FOR THE PREVENTION OF FLY STRIKE AFTER MULESING (<i>Diagram 2</i>)					<i>Diagram 3 (Area of sheep to treat after marking)</i>
Apply two bands, one over each side of the mulesing wound and overlapping 25 mm onto the surrounding wool.					
SHEEP WEIGHT (kg)	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	NO. OF SHEEP TREATED PER [1 L][5 L] [10 L][15 L][20 L][22 L]	8. This product is not recommended for treating existing strikes. Use a product registered as a dressing for existing strikes. 9. Heavy rain following the application of this product could diminish the period of protection. 10. STRIKEFORCE-S Spray-On Sheep Blowfly Treatment contains a scourable colouring agent for easy band identification. 11. Withholding periods must be observed following application of STRIKEFORCE-S Spray-On Sheep Blowfly Treatment for any purpose including after mulesing.
Up to 30	8	4	2	[125][625][1250] [1875][2000][2750]	
31 - 50	10	5	2	[100][500][1000] [1500][2000][2200]	
FOR THE PREVENTION OF FLY STRIKE AFTER MARKING (<i>Diagram 3</i>)					
<ul style="list-style-type: none"> When a tail has been amputated, apply 4 mL over the crutch, ensuring sufficient coverage of the open wound. When the tail has not been amputated, ensure sufficient coverage of the tail where the rubber ring has been applied. In the case of ram lambs a further 4 mL can be sprayed over and around the scrotum area after the rubber ring has been applied.					
LAMB WEIGHT (30 kg)	TOTAL VOLUME / LAMB (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	AVERAGE NO. OF LAMBS TREATED PER [1 L][5 L][10 L] [15 L][20 L][22 L]	
Ram lambs	8	4	2	[166][833][1666]	
Ewe lambs	4	4	1	[2500][3333][3666]	