



Product Name: Caboose 500SC Fungicide
APVMA Approval No: 82417/112107

Label Name:	Caboose 500SC Fungicide
Signal Headings:	DANGEROUS POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT: 500g/L CARBENDAZIM
Mode of Action:	GROUP 1 FUNGICIDE
Statement of Claims:	A broad spectrum, systemic fungicide for the control of fungal diseases in various crops as per the Directions for Use table.
Net Contents:	1L, 5L, 10L 20L
Restrains:	DO NOT apply to turf, grapes, cucurbits (including melons), citrus fruit, custard apple, mango, pome fruit (apples or pears) or stone fruit (including cherries).
Directions for Use:	
Other Limitations:	WARNING: Contains carbendazim which causes birth defects and irreversible male infertility in laboratory animals. Avoid contact with carbendazim
Withholding Periods:	MACADAMIA NUTS: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION

	<p>DO NOT ALLOW LIVESTOCK TO GRAZE VEGETATION IN TREATED PLANTATIONS FOR 4 WEEKS AFTER APPLICATION. DO NOT CUT TREATED VEGETATION IN TREATED PLANTATIONS FOR STOCK FEED FOR 4 WEEKS AFTER APPLICATION</p> <p>CHICKPEAS, FABA BEANS, LENTILS AND VETCH: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION DO NOT GRAZE OR CUT FOR STOCKFEED FOR 4 WEEKS AFTER APPLICATION</p>
--	---

Trade Advice:	<p>EXPORT OF TREATED PRODUCE Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Caboose 500SC Fungicide. If you are growing produce for export, please check with Relyon (Australia) Pty Ltd for the latest information on MRLs and import tolerances BEFORE using Caboose 500SC Fungicide.</p>
----------------------	--

General Instructions:	<p>MIXING / APPLICATION SHAKE WELL BEFORE USE. Caboose 500SC Fungicide is a liquid suspension to be mixed with water for application as a spray. Add the required quantity of Caboose 500SC Fungicide to a partly filled spray tank and agitate. Complete filling while agitating. In the event of a stoppage during spraying, ensure the spray mixture is properly agitated before spraying recommences.</p> <p>ADDITION OF WETTING AGENT The addition of a wetting agent e.g. Activator Surfactant to spray mixture improves the distribution of the spray on hard to wet plants.</p>
------------------------------	---

Resistance Warning:	<p>GROUP 1 FUNGICIDE</p> <p>Caboose 500SC Fungicide is a member of the benzimidazole group of fungicides. For fungicide resistance management Caboose 500SC Fungicide is a Group 1 fungicide. Some naturally occurring fungal populations resistant to Caboose 500SC Fungicide and other Group 1 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Caboose 500SC Fungicide and other Group 1 Fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Relyon (Australia) Pty Ltd accepts no liability for any losses that may result from the failure of Caboose 500SC Fungicide to control resistant fungi.</p>
----------------------------	--

Precautions:	<p>PRECAUTION Keep away from foodstuffs.</p> <p>Re-entry Period DO NOT allow entry into treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.</p>
---------------------	--

Protections:	<p>PROTECTION OF LIVESTOCK Caboose 500SC Fungicide is of low toxicity to bees.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Very toxic to aquatic life with long lasting effects. DO NOT contaminate dams, rivers or waterways with the chemical or used containers.</p> <p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS</p>
---------------------	--

DO NOT apply under weather conditions, or from spraying equipment that may cause spray or drift onto into nearby susceptible plants / crops, cropping lands or pastures.

Storage and Disposal:

STORAGE AND DISPOSAL

Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilizers.

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or preferably pressure rinse containers before disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection point or similar container management program site. The cap should not be replaced, but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers OR product.

Safety Directions:

Very Dangerous, particularly the concentrate. Poisonous if absorbed by skin contact, inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. When opening the container, preparing spray and using the prepared spray wear elbow-length chemical resistant gloves and a face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

First Aid Instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26).

First Aid Warnings:

DIRECTIONS FOR USE

TREE CROPS

RATE					CRITICAL COMMENTS
This table shows rates for dilute spraying. For concentrate spraying, refer to mixing/application section					For all uses in this table: Apply by dilute or concentrate spraying application equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Macadamia nuts	Macadamia Husk Spot (<i>Pseudocercospora macadamiae</i>)	All States	50mL plus a wetting agent (at 100mL/100 L)	14 days (H) 4 weeks (G)	Apply at 5 and 8 weeks after main flowering – Stage 2 anthesis (white flowering stage). Remove any fallen nuts from under trees prior to spraying. DO NOT apply more than 2 applications of Caboose 500SC Fungicide per season.

OTHER CROPS

Crops	Disease	State	Rate	WHP	Critical Comments
Chickpeas, faba beans, lentils and vetch	Chocolate spot, (<i>Botrytis fabae</i>), Grey Mould (<i>Botrytis cinerea</i>)	All States	500mL/ha	4 weeks (H,G)	Apply a maximum of two consecutive applications at 14 day intervals. Apply in a minimum of 100L water per hectare.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.