

Product Name: Conquest Imazamax 700 WG Herbicide
APVMA Approval No: 83146/133682



Label Name:	Conquest Imazamax 700 WG Herbicide
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	700 g/kg IMAZAMOX
Mode of Action:	GROUP 2 HERBICIDE
Statement of Claims:	For the post-emergence control of certain annual grass and broad leaved weeds in Clearfield Plus wheat, Clearfield barley, field peas, legume-based pastures, lucerne, peanuts and soybeans, as specified in the Directions for Use table.
Net Contents:	0.5 kg - 20 kg
Restrains:	

Directions for Use:	
---------------------	--

Other Limitations:	
--------------------	--

Withholding Periods:	<p>GRAZING Lucerne, Pastures: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION. Peanuts, Soybeans: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION. Field Peas: DO NOT GRAZE OF CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION. CLEARFIELD PLUS WHEAT AND CLEARFIELD BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION. HARVEST Field peas, Peanuts, Soybeans: NOT REQUIRED WHEN USED AS DIRECTED.</p>
----------------------	---

Trade Advice:	
---------------	--

General Instructions:	
-----------------------	--

Resistance Warning:	<p>Imazamax 700 WG Herbicide is a member of the imidazoline group of herbicides. This product has the ALS mode of action. For weed resistance management Imazamax 700 WG Herbicide is a Group 2 Herbicide. Some naturally occurring weed biotypes resistant to the product and other Group 2 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 2 Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Crop Protection Pty Ltd accepts no liability for any losses that may result from failure of this product to control resistant weeds.</p>
---------------------	---

Precautions:	<p>RE-ENTRY DO NOT re-enter treated areas until spray has dried. If re-entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves.</p>
--------------	---

Protections:

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Field Peas

DO NOT apply to field peas beyond the 4 node stage.

DO NOT apply if frost is forecast.

DO NOT use Imazamax 700 WG Herbicide after simazine (unregistered in field peas) and DO NOT exceed 250g ac/ha diuron if use of Imazamax 700 WG Herbicide is planned. DO NOT roll field peas after germination if use of Imazamax 700 WG Herbicide is planned.

Before using this product on field pea varieties other than Bohatyr, Bonzer, Dun, Dundale, Glenroy, Laura, Magnet, Mukta, Santi and Wirrega, contact Conquest Crop Protection Pty Ltd for up-to-date information.

DO NOT use on Alma, Excell and Parafield varieties.

Pasture Legumes and Perennial Grasses

Safety to medic varieties may be lower than to clover varieties. Safety to seedling pastures less than once year old may be lower than to older established pastures. Transient biomass reduction and leaf yellowing may occur to pasture legumes and perennial grass, with good recovery occurring under favourable seasonal conditions. In some situations a reduction in perennial grass numbers may occur as a result of unfavourable seasonal conditions following application. For up-to-date information on all pasture clover and medic varieties and other temperate pasture species, contact Conquest Crop Protection Pty Ltd before use.

Lucerne

Before using this product on lucerne varieties other than Hunterfield, Sceptre, Sequel, Siriver and Trifecta, contact Conquest Crop Protection Pty Ltd for up-to-date information.

DO NOT USE THE GLISARIN® MIX ON SEEDLING LUCERNE.

Replanting

If replanting is necessary in a field previously treated with Imazamax 700 WG Herbicide, the field may be replanted to field peas (above listed varieties only), soybeans or peanuts. DO NOT apply a second treatment of Imazamax 700 WG Herbicide.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to aquatic flora. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

Storage and Disposal:	<p>STORAGE AND DISPOSAL</p> <p>Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.</p>
-----------------------	---

Safety Directions:	<p>Will irritate the eyes. When opening the container and preparing spray, wear face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash face shield or goggles.</p>
--------------------	--

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE

Crop use or Situation	Weeds controlled	Rate	Critical Comments
<p>Clearfield Plus wheat DO NOT use on CL STL and CL JNZ wheat varieties.</p> <p>Clearfield barley</p>	<p>Brome (<i>Bromus diandrus</i> and <i>B. rigidus</i>) Barley (<i>Hordeum vulgare</i>) Barley grass (<i>Hordeum leporinum</i>) Indian hedge mustard (<i>Sisymbrium orientale</i>) Muskweed (<i>Myagrum perfoliatum</i>) Oat (<i>Avena sativa</i>) Triticale (<i>Triticosecale spp</i>) Wheat (<i>Triticum aestivum</i>) – non Clearfield* varieties Wild oat (<i>Avena fatua</i>) Wild radish (<i>Raphanus raphanistrum</i>) Wild turnip (<i>Brassica tournefortii</i>)</p>	<p>17.7 – 35.4 g/ha tank mix with 7.5 - 15 g/ha Conquest Gunner Herbicide, plus Glisarin®, or Hasten* at 0.5L/100L spray volume</p>	<p>NOT for use on Clearfield* wheat varieties Clearfield* STL and Clearfield* JNZ.</p> <p>Apply to Clearfield Plus wheat crops from the 3 leaf stage to 1st node stage (Z31). Apply to Clearfield barley crops from the 5 leaf stage to 1st node stage (Z31).</p> <p>DO NOT apply to Clearfield Plus wheat or Clearfield barley after the 1st node stage (Z31).</p> <p>Applications should be targeted at grass weeds when the majority are in the 2-4 leaf stage and only when within the recommended crop stages. Application to multi-tillered crops may impair weed control because of poor contact and coverage of weeds. – see Compatibility. Tank mixes with LontrelTM Advanced Herbicide at 75mL/ha will provide control of composite and legume weeds. Tank mixes with Conquest 570 LVE MCPA Herbicide at 500mL/ha will provide control of composite and brassicaceous weeds. The control of brassicaceous weeds will depend on the status of Group 2 resistance in the population. The addition of Conquest 570 LVE MCPA will improve control and provide an additional mode of action for resistance management. If other weeds require control, apply appropriate herbicides at least two weeks before or after Imazamax and Gunner tank mix and only when signs of regrowth or renewed vigour appear, otherwise the effects of the early treatment may affect the performance of the subsequent treatment.</p>
	<p>Annual ryegrass (<i>Lolium rigidum</i>) Dense flower fumitory (<i>Fumaria densiflora</i>) Marshmallow (<i>Malva parviflora</i>) Subclover (<i>Trifolium subterraneum</i>) Suppression Bedstraw spp. (<i>Galium tricornutum</i> and <i>G. aparine</i>) Doublegee (<i>Emex australis</i>) Silver grasses (<i>Vulpia bromoides</i> and <i>V. myuros</i>)</p>	<p>28.3 – 35.4 g/ha plus 12 - 15 g/ha Conquest Gunner Herbicide, plus Glisarin®, or Hasten* at 0.5L/100L spray volume</p>	<p>Weed species will either be controlled or suppressed. In both cases, surviving plants will be stunted and will be uncompetitive with the crop, and seed set will be prevented or greatly reduced. The control of annual ryegrass varies from excellent to poor depending on the status of Group 2 resistance in the population and environmental conditions. Where the population is expected to exceed 200 plants/m², or a high level of control is required, or the ryegrass is known to be resistant or thought to be developing resistance, an application of a suitable pre-emergent should be made prior to sowing. A follow up grass selective herbicide may also be necessary.</p>

Important:

- Before using Imazamax in tank mixes with other herbicides, check the plant-back information on all product labels. The time between spraying and planting will be determined by the product with the longest plant-back period.

CROP	WEEDS CONTROLLED	RATE/Ha	WHP	CRITICAL COMMENTS
Field Peas Post-emergence See "Crop Safety" re varieties	Barley grass (<i>Hordeum leporinum</i>), Brome grass (<i>Bromus</i> spp.), Deadnettle (<i>Lamium amplexicaule</i>), Indian Hedge mustard (<i>Sisymbrium orientale</i>), Storksbill (<i>Erodium</i> spp.), Turnip weed (<i>Rapistrum rugosum</i>), Volunteer barley (<i>Hordeum vulgare</i>), Volunteer oats (<i>Avena sativa</i>), Volunteer lupins (<i>Lupinus</i> spp.), Volunteer triticale (<i>Triticosecale</i> spp.), Volunteer wheat (<i>Triticum aestivum</i>), Wild Oats (<i>Avena</i> spp.), Wild Turnip (<i>Brassica tournefortii</i>), Doublegee (<i>Emex australis</i>), *Shepherds purse (<i>Capsella bursa-pastoris</i>), * Threehorn bedstraw (<i>Calium tricomutum</i>), *Wild radish (<i>Raphanus raphanistrum</i>), *Wireweed (<i>Polygonum aviculare</i>)	45g plus Wetter 1000, or equivalent, at 200mL per 100L water	6 weeks (grazing)	<p>DO NOT apply to field peas beyond the 4-node stage. DO NOT use adjuvants other than Wetter 1000 or equivalent. Apply to actively growing broadleaved weeds in the cotyledon to 3-leaf stage. Apply to grass weeds up to the 2-tiller stage. Good crop growth with aid weed control. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. *Control will not be 100% but surviving plants will generally be retarded and will not complete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds. Leaf yellowing and height suppression may occur, especially under adverse growing conditions such as moisture stress. Flowering may be delayed.</p> <p>IMPORTANT: Refer to CROP SAFETY section of label re use on field peas.</p> <p>Refer to FOLLOW CROPS section of this label regarding follow crops.</p>
Legume- based pastures Post-emergence Clovers, Lucerne, Medics, Serradellas only; see "Crop Safety" re varieties		<p>45g plus Glisarin® at 500mL per 100L water</p> <p>50g plus Glisarin® at 500mL per 100L water plus Amsul 417 at 2L per 100L water</p>	7 days (grazing)	<p>Seedling - Apply to legumes after full emergence of the third trifoliolate leaf but well before commencement of flowering. Established- Apply before commencement of flowering. Apply to actively growing broadleaved weeds in the cotyledon to 3-leaf stage. Apply to grass weeds up to the 2-tiller stge. Good pasture growth will aid weed control. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Where pasture stand is poor, weed control may be inadequate due to lack of competition. Ensure sufficient spray coverage of weeds particularly in dense pastures. *Control will not be 100% but surviving plants will generally be retarded and will not complete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds. Refer to CROP SAFETY section of this label re varieties of clovers and medics and other pasture species. Note: use of the 50g/ha rate plus Glisarin® may severely damage medics. Refer to FOLLOW CROPS section of this label regarding follow crops.</p>

CROP	WEEDS CONTROLLED	RATE/Ha	WHP	CRITICAL COMMENTS
Lucerne Post-emergence (See "Crop Safety" re varieties)	As for post emergence use in field peas (winter weeds) and soybeans (summer weeds)	45 or 50g plus WETTER 1000, or equivalent, at 200mL per 100L water plus AMSUL 417 at 2L per 100L water	7 days (grazing)	Apply to actively growing weeds at growth stages as for field peas (winter weeds) and soybeans (summer weeds). DO NOT use the Hasten or Kwicken mix on seedling lucerne. Use the 50g rate for summer weeds. Crop Stage: Seedling: Apply when crop is at 2 trifoliate leaf stage and before commencement of flowering. Established: Apply as above following cutting or grazing. Refer to CROP SAFETY section of this label re varieties of lucerne
Post-emergence Peanuts, Soybeans	Amaranth (<i>Amaranthus</i> spp.), Barley grass (<i>Hordeum leporinum</i>), Barnyard grass (<i>Echinochloa crus-gali</i>), Bellvine (<i>Ipomoea plebeia</i>), Brome grass (<i>Bromus</i> spp.), Deadnettle (<i>Lamium amplexicaule</i>), Fathen (<i>Chenopodium album</i>), Fierce Thornapple (<i>Datura ferox</i>), Indian Hedge mustard (<i>Sisymbrium orientale</i>), Liverseed grass (<i>Urochloa panicoides</i>), Turnip weed (<i>Rapistrum rugosum</i>), Volunteer barley (<i>Hordeum vulgare</i>), Volunteer oats (<i>Avena sativa</i>), Volunteer lupins (<i>Lupinus</i> spp.), Volunteer triticale (<i>Triticosecale</i> spp.), Volunteer wheat (<i>Triticum aestivum</i>), Wild gooseberry (<i>Physalis minima</i>), Wild Oats (<i>Avena</i> spp.), Wild Turnip (<i>Brassica tournefortii</i>), *Anoda weed (<i>Anoda cristata</i>), *Awnless barnyard grass (<i>Echinochloa colona</i>), *Blackberry nightshade (<i>Solanum nigrum</i>), *Caltrop (<i>Tribulus terrestris</i>), *Chickweed (<i>Stellaria media</i>), *Crabgrass (<i>Digitaria ciliaris</i>), *Doublegee (<i>Emex australis</i>), *Noogoora burr (<i>Xanthium pungens</i>), *Shepherds purse (<i>Capsella bursa-pastoris</i>), * Threehorn bedstraw (<i>Calium tricomutum</i>), *Wild radish (<i>Raphanus raphanistrum</i>), *Wireweed (<i>Polygonum aviculare</i>)	50g plus EITHER WETTER 1000 (or equivalent), at 200mL/100L water plus AMSUL 417 at 2L/100L water OR Glisarin®, at 500mL or 1L /100L water	4 weeks (grazing)	DO NOT apply to crops beyond the 6-leaf stage. Apply to actively growing weeds in the cotyledon to 2- leaf stage. Apply to grass weeds up to the 2-tiller stage. If using the Glisarin® mix, use the 1L/100L rate where grass weeds predominate. *Control will not be 100% but surviving plants will generally be retarded and will not compete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds. Refer to FOLLOW CROPS section this label regarding follow crops.

* Registered trademarks

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

This product is for early post-emergence weed control in field peas, lucerne, legume-based pastures, soybeans and peanuts. In addition, this product is safe for use in Clearfield Plus wheat and Clearfield barley (as per above use pattern). Weeds will either die or remain stunted and will not compete with the crop. For weeds marked with an asterisk in the above table, the level of control will vary significantly depending on factors such as climatic conditions following application, crop vigour, weed seed depth, etc.

A spray adjuvant **MUST** always be added, as per the "Directions for Use" table, while Boost liquid ammonium sulphate should also be added in situations indicated in the table.

Good crop competition is essential for effective weed control.

MIXING

Imazamax 700 WG Herbicide is a water dispersible granular formulation. Part fill the spray tank with water, then with the agitator running, add the required amount of product, then fill the tank with water. When tank mixing with other products, follow the industry recommended mixing order by formulation types. For more complex mixtures contact a Conquest representative.

DO NOT tank mix with either MCPA Amine or 2,4-D Amine when using in pastures

For field peas, lucerne, legume-based pastures, soybeans and peanuts: DO NOT tank mix with selective post-emergence grass herbicides. DO NOT apply these herbicides following use of Conquest Imazamax 700 WG Herbicide until grasses have resumed active growth. Alternatively, apply the grass herbicide at least one day before using Conquest Imazamax 700 WG Herbicide.

APPLICATION

For ground application only. Apply with boom equipment in not less than 50L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands.

DO NOT apply by aircraft.

DO NOT apply by mister.

Imazamax 700 WG Herbicide should be applied a minimum of two hours before rainfall or irrigation.

DO NOT apply Imazamax 700 WG Herbicide more than once per growing season.

EQUIPMENT CLEANUP

Thoroughly flush all spray equipment with water following the use of Imazamax 700 WG Herbicide and before use with other products. Rinse water should NOT be discharged where it will reach streams, water bodies or natural vegetation.

COMPATIBILITY

Imazamax is chemically compatible with Alpha Forte and Alpha Duo Insecticides. It is also chemically and biologically compatible with Conquest Gunner Herbicide, Conquest Nitro 240 Herbicide, Conquest Recon 520 Herbicide, Conquest Corsair, Conquest 570 LVE MCPA, Conquest 750 MCPA Herbicide, Conquest Bromo 200 and Conquest Brom M Herbicides, Conquest Racer and Conquest Crossfire Herbicides. For the use with other products contact a Conquest representative.

Mixes with Lontrel^{TM5} Advanced in Clearfield Plus wheat and Clearfield barley

Tank mixes with Lontrel^{TM5} Advanced at 150mL/ha will provide control of composite and legume weeds. Refer to Lontrel^{TM5} Advanced label.

Mixes with Conquest 570 LVE Herbicide in Clearfield Plus wheat and Clearfield barley

Tank mixes with Conquest 570 LVE Herbicide at 500mL/ha will provide control of composite and brassicaceous weeds. Refer to Conquest 570 LVE label.

FOLLOW CROPS

Before using Imazamax in tank mixes with other herbicides, **check plant-back information on all product labels**. The time between spraying and planting will be determined by the product with the longest plant-back period.

Under conditions, such as very dry season, which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible. The following minimum re-cropping intervals (months after application) should be observed.

Following use in lucerne, legume-based pastures and field peas:

Months after application		
0	10	21
Field peas	Chickpeas	All other crops including canola (except varieties with CLEARFIELD technology) Oats Safflower
Canola varieties with CLEARFIELD technology	Faba Beans	
Maize varieties with CLEARFIELD technology	Lucerne	
Wheat varieties with CLEARFIELD technology	Lupins	
	Pasture Legumes	
	Vetch	
	*Barley	
	*Wheat (except varieties with CLEARFIELD technology)	
	*Triticale	

*The following additional requirements apply if it is intended to sow WHEAT, BARLEY or TRITICALE during the next winter season:

DO NOT apply Imazamax 700 WG Herbicide later than the end of August.

DO NOT use Imazamax 700 WG Herbicide in areas where rainfall from time of spraying to sowing of cereals is expected to be below 200mm.

Furthermore:

DO NOT use on soils of pH 5.5 (CaCl₂) or less in areas where rainfall from spraying to sowing of cereals is expected to be below 300mm.

In late, short seasons where the soil is cold for most of the time it is wet, break-down will be slower and plant-back times will be extended.

If expected rainfall is not received following use of Imazamax 700 WG Herbicide, consult your local Conquest Crop Protection representative before planting wheat, barley or triticale. (In calculating rainfall actually received, exclude single, isolated falls which do not result in periods of continuous soil moisture to allow microbial breakdown to occur in the root zone).

Following use in summer crops.

Irrigated only

Months after application			
0	5	10	18
Mung beans Peanuts Soybeans Canola varieties with CLEARFIELD technology Maize varieties with CLEARFIELD technology Wheat varieties with CLEARFIELD technology	*Chickpeas *Lucerne *Lupins *Pasture Legumes *Barley *Wheat (except varieties with CLEARFIELD technology)	**Maize (except varieties with CLEARFIELD technology) **Sorghum **Cotton **Oats **Sunflower	All other crops (providing rainfall and irrigation exceeds 2000mm)

*DO NOT plant these crops unless interim rainfall (rainfall plus irrigation) from application to sowing is at least 500mm.

** DO NOT plant these crops unless interim rainfall (rainfall plus irrigation) from application to sowing is at least 800mm.

Dryland only

Months after application				
0	8	10	15	22
Mung beans Peanuts Soybeans Canola varieties with CLEARFIELD technology Maize varieties with CLEARFIELD technology Wheat varieties with CLEARFIELD technology	*Lucerne *Barley *Wheat (except varieties with CLEARFIELD technology)	**Chickpeas **Maize (except varieties with CLEARFIELD technology) **Sorghum **Cotton **Oats **Sunflower	Pasture Legumes Lupins	All other crops (providing rainfall and irrigation exceeds 2000mm)

*DO NOT plant these crops unless interim rainfall (rainfall plus irrigation) from application to sowing is at least 500mm.

**DO NOT plant these crops unless interim rainfall (rainfall plus irrigation) from application to sowing is at least 800mm.

When Tank- Mixed with Conquest Gunner 750

The following minimum re-cropping intervals (months after application) should be observed.

Months after application		
0	10	34
Clearfield Plus Wheat Clearfield Wheat Clearfield barley Clearfield canola	chick peas, faba beans, field peas, lucerne, lupins, pasture legumes, *oats *triticale Vetch, non-Clearfield Barley, Non-Clearfield Wheat	Conventional and other herbicide tolerant canola, All other crops.

*Non-Clearfield Barley, Oats, Triticale, Non-Clearfield Wheat:

The following additional requirements apply if it is intended to sow these cereals during the next winter season:

- DO NOT apply Imazamax and Gunner later than the end of August in winter crops (no later than the end of July in WA).
- DO NOT use Imazamax and Gunner in areas where rainfall from spraying to sowing of cereals is expected to be below 150 mm (for respectively, 18 and 7.5 g/ha use), 200 mm (for respectively up to 23.5 and 10 g/ha use) and 250 mm (for respectively 28.3

and 12 g/ha – 35.3 and 15 g/ha use).

- DO NOT use above 17.7 and 7.5 g/ha in the Lower Great Southern region of Western Australia.

For all situations, if expected rainfall is not received following use of Imazamax and Gunner, consult your local Conquest representative before planting non-Clearfield cereals. In calculating rainfall actually received, place greater emphasis on rain received from application up to the end of Spring and lesser emphasis on break rains. If single isolated heavy summer and autumn falls and break rains are required to achieve rainfall targets, it may not be safe to sow non-Clearfield cereals within 10 months of application. Consult your local Conquest representative for advice.

CROP SAFETY

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth of Clearfield Plus wheat and Clearfield barley, however plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under poor growth conditions.

The **Clearfield** wheat varieties **Clearfield** STL and **Clearfield** JNZ have limited tolerance to **Imazamax** Herbicide and **Gunner** tankmix combination. Application of **Imazamax and Gunner tank mix combination** to these wheat varieties can cause unacceptable injury especially under cold, wet conditions. **Imazamax** Herbicide plus gunner should not be used on these varieties.

This product may cause shortening of plant internodes and may, in some circumstances, lead to transient crop yellowing but plants will soon recover. This effect is more pronounced under poor growth conditions including conditions of prolonged moisture stress.

Restrictions:

- Apply ONLY to certified Clearfield Plus wheat and Clearfield barley, varieties with the Clearfield technology, excluding Clearfield STL and Clearfield JNZ. DO NOT apply to conventional or other herbicide tolerant wheat and barley varieties.
- DO NOT apply to crops that are stressed due to conditions such as waterlogging, too little moisture, frost, disease or nutritional disorders.
- DO NOT apply by aircraft.
- DO NOT apply if rain is expected within 2 hours of application. DO NOT apply more than once per season to any one crop.
- DO NOT use in Clearfield Plus wheat crops in tank mix or sequentially with diuron, sulfonamides, or sulfonamides.

Spray Drift Restrictions:

Specific definition of the terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystander to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from those sensitive areas.

DO NOT apply unless wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evening one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplet smaller than a MEDIUM spray droplet size category.
- Minimum distance between the application site and downwind sensitive area (see 'Mandatory buffer zones' section of the following tables titled 'Buffer zones for boom sprayers') are observed.

Buffer zone for boom sprayers when applied alone

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 50 g /ha	0.5 m or lower	0 metres	5 metres	0 metres	10 metres	0 metres
	1.0 m or lower	0 metres	30 metres	0 metres	45 metres	0 metres

Buffer zone for boom sprayers when applied in tank mix combination with CONQUEST GUNNER 750 WG HERBICIDE

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 35.4g/ha Conquest Imazamax 700 WG Herbicide + 15 g/ha Conquest Gunner 750 WG Herbicide	0.5 m or lower	0 metres	0 metres	0 metres	25 metres	0 metres
	1.0 m or lower	0 metres	20 metres	0 metres	70 metres	0 metres