

Product Name: FENNEC POUR-ON FOR SHEEP
APVMA Approval No: 85544/123481



Label Name:	FENNEC POUR-ON FOR SHEEP
-------------	--------------------------

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	35 mg/ml IMIDACLOPRID 4 mg/ml ABAMECTIN Solvent: 300 mg/ml N-METHYL-2-PYRROLIDONE
-------------------------	---

Claims:	<p>For the treatment and control of susceptible strains of lice (<i>Bovicola ovis</i>) on sheep, including synthetic pyrethroid resistant lice, when applied up to 7 days after shearing. Sheep will be protected against re-infestation with lice for 5 weeks after treatment when applied within 24 hours of shearing.</p> <p>For the treatment and control of abamectin susceptible Small Brown Stomach Worm, Black Scour Worm, Stomach Hair Worm, Nodule worm and Large Bowel worm.</p> <p>Fennec Pour-On for Sheep is effective for the treatment and control of the following abamectin sensitive adult and immature gastrointestinal nematodes: Small Brown Stomach Worm (<i>Teladorsagia circumcincta</i>) Black Scour Worm (<i>Trichostrongylus</i> spp.) Stomach Hair Worm (<i>Trichostrongylus axei</i>) Nodule Worm (<i>Oesophagostomum columbianum</i>) Large Bowel Worm (<i>Oesophagostomum venulosum</i>) Fennec Pour-On for Sheep contains imidacloprid, a member of the neonicotinoid family of chemicals and abamectin, a member of the macrocyclic lactone family of chemicals. Resistance may develop to any chemical.</p>
---------	--

Net Contents:	10 L 100 L 110 L 120 L 15 L 20 L 22 L 25 L 5 L 55 L
---------------	--

	60 L
--	------

Directions for Use:	
---------------------	--

Restrains:	DO NOT USE on ewes which are producing, or may in the future produce milk that may be used or processed for human consumption.
------------	--

Contraindications:	<p>DO NOT USE on unshorn sheep, or shorn sheep more than 7 days after shearing.</p> <p>DO NOT USE on ewes with unshorn lambs at foot.</p> <p>This product should not be used on lambs under 6 weeks of age or less than 10 kg bodyweight.</p> <p>This product should not be used on severely sick, debilitated or dying animals.</p> <p>This product should not be used on sheep which have not been cleanly shorn for any reason e.g. Mycotic Dermatitis.</p>
--------------------	--

Precautions:	This product is for application as a pour-on only. Do not give orally or by injection. Do not apply this product at the same time as any other pour-on products.
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	<p>Dose rate: 10 mL for each 10 kg body weight.</p> <p>Sheep should be weighed before treatment.</p> <p>Apply the entire dose to sheep under 30kg as a single broadband stripe from poll of the head to the base of the tail.</p> <p>For sheep over 30kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the base of the tail. The stripes should be no further than 10 cm apart. Refer to diagrams below.</p> <p>The following table may be used as a guide:</p> <p>Animals heavier than 90 kg to be dosed at 10 mL per 10 kg. A representative sample of animals should be weighed before treatment. Dose the mob according to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.</p>
----------------------------	---

General Directions:	<p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.</p> <p>Apply product only with the recommended applicator fitted with a six-hole T-bar nozzle. Check the accuracy of equipment before and during treatment. Efficacy is not affected by heavy rainfall before and/or after treatment.</p> <p>Avoid treating ewes less than 6 weeks before lambing commences, while ewes are lambing or have lambs at foot, as live lice can persist and infect the lambs.</p> <p>Treated sheep should not be mixed with lice free sheep until 6 weeks after treatment.</p> <p>Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time, no split shearings, stock-proof fencing to prevent external introductions, quarantine and lice treatment of all newly purchased stock are essential for effective lice control.</p>
---------------------	--

	<p>Fennec Pour-On for Sheep contains a blue dye to identify treated sheep. This dye is scourable from wool.</p> <p>Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted before any parasite treatment is used.</p> <p>Do not leave container, hose or applicator in the sun for long periods of time.</p> <p>The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces.</p>
--	---

Withholding Periods:	<p>MEAT: DO NOT USE less than 63 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE on ewes which are producing, or may in the future produce milk that may be used or processed for human consumption.</p> <p>WOOL: DO NOT USE less than 6 months before shearing or fibre collection.</p> <p>Use of this product may result in wool residues that may not comply with European Union environmental quality standards.</p> <p>Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.</p>
----------------------	---

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 84 days before slaughter for export. Before using this product, confirm the current ESI from Boehringer Ingelheim on 1800 808 691 or the APVMA website (www.apvma.gov.au/residues)</p>
---------------	--

Safety Directions:	<p>Harmful if swallowed. Will irritate the eyes. Avoid contact with eyes. When using the product wear cotton overalls, buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield or goggles. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.</p> <p>Rehandling Interval: DO NOT handle sheep after treatment until dry.</p>
--------------------	--

First Aid Instructions:	<p>If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. If in eyes wash out immediately with water.</p>
-------------------------	--

First Aid Warnings:	
---------------------	--



Additional User Safety:	<p>Additional information is listed in the Safety Data Sheet.</p>
-------------------------	---

Environmental Statements:	<p>ENVIRONMENTAL PROTECTION</p> <p>Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.</p>
---------------------------	--

Disposal:	<p>This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple rinse or pressure rinse container before disposal. Dispose of rinsate or any undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.</p>
-----------	--

Storage:

Store below 30°C (Room Temperature). Protect from light.

Bodyweight (kg)	Dose Volume (mL)	Method	Apply in the pattern below
10 to 12.5	12.5	Single stripe	
12.6 to 15	15		
15.1 to 20	20		
20.1 to 25	25		
25.1 to 30	30		
30.1 to 40	40	Double stripe	
40.1 to 50	50		
50.1 to 60	60		
60.1 to 70	70		
70.1 to 80	80		
80.1 to 90	90		