

Product Name: Conquest In2 Pro Crop Oil Concentrate Spray Adjuvant

APVMA Approval No: 85618/113756

Label Name:	Conquest In2 Pro Crop Oil Concentrate Spray Adjuvant		
Signal Headings:	CAUTION		
	KEEP OUT OF REACH OF CHILDREN		
	READ SAFETY DIRECTIONS BEFORE OPENING OR USING		
Constituent	Actives: 582 g/L Paraffinic oil		
Statements:	240 g/L Nonionic surfactants		
Mode of Action:			
Statement of Claims:	A crop oil concentrate spray adjuvant to improve the spreading and wetting of herbicides, insecticides and defoliants on plant surfaces as per directions for use.		
	inconstant and actionalities on plant carriades as per uncontaine for acci.		
Net Contents:	20-1000L		
Net Contents.	20-1000L		
Restraints:			
Directions for Use:			
Other Limitations:			
Withholding Periods:	Adhere to the withholding period(s) specified on the label of the companion product(s) used.		
Trade Advice:			

#### General Instructions:

#### **MIXING**

These are general mixing instructions. Always follow the mixing instructions of each ingredient in the tank-mix. Ensure product compatibility. Recommendations on other product labels take precedence over mixing instructions on this label.

- Add water to the spray tank to 10cm above the level of agitation and ensure the agitation device is working vigorously (there must be a minimum of 100L of water in the tank before any pesticide is added).
- If tank mixing, use the following general order for mixing partners. First add water dispersible granules (e.g. Eclipse), second add flowable formulations (e.g. Atrazine), third add emulsifiable concentrates (e.g. Starane\*200EC), fourth add water-soluble liquids (e.g. Lontrel\* Herbicide) and last add In2 Pro and other adjuvants.
- Between the additions of each ingredient allow at least one minute to ensure thorough mixing before the new tank-mix partner is added.
- Finally add In2 Pro. Continue filling the tank to the required volume maintaining agitation at all times.
- Only mix sufficient spray solution for immediate use. Verdict\* 520 and any other tankmixes should be applied immediately for best results.

#### STORAGE OF SPRAY SOLUTION

Agitate for 1 hour every 8 hours, and agitate for 1 hour before spraying out where spray solution is not applied within 8 hours of mixing.

# **CLEANING SPRAY EQUIPMENT**

The presence of product(s) makes it essential for the spray equipment to be cleaned thoroughly between different applications. To rinse and decontaminate spray equipment refer to relevant product labels for instructions.

#### Resistance Warning:

### Precautions:

Explosion Warning: Apparently 'Empty' containers may contain liquid and/or vapour, which can be explosive if exposed to an ignition source at temperatures above 90oC. Such conditions may occur during cutting or welding. DO NOT cut or weld these containers. Explosion may cause injury or death.

## Protections:

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with the chemical or used container.

# Storage and Disposal:

Store in the closed original, container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. DO NOT store near food, feedstuffs, fertiliser or seed. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Do not burn empty containers or product.

Recyclable Containers (110L): Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### SMALL SPILL MANAGEMENT

Wear appropriate clothing whilst cleaning up small spills. Apply absorbent material such as earth, cat litter, sand or clay granules to the spill. Sweep up material for disposal when absorption is complete. Dispose of the contaminated material in accordance with local and/ or State recommendations.

Safety Directions:	Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When opening container, preparing spray wear elbow-length PVC gloves and face shield or goggles. If product gets in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles and contaminated clothing. When using with another chemical, consult the safety directions of the companion product.
First Aid Instructions:	If poisoning occurs, contact a doctor of Poisons Information Centre. Phone Australia 13 11 26.
First Aid Warnings:	

#### DIRECTIONS FOR USE

**Important:** Prior to use read all directions on this label and on the label of the companion crop protection product. The product(s) label for the specific usage and restrictions have precedence over Directions for Use referred to on this label.

In 2 Pro is compatible with a wide range of herbicides including:

Atrazine (Atraquest 900 WG) X-pand\* Herbicide Axial\* Herbicde Starane\* Advance Herbicide Nitro® 240 EC Herbicide Racer® 250 WG Maca® 600 Neon® 400 Velocity\* Herbicide Simazine (Simaguest 900 WG) Broadstrike \* Herbicide Eclipse\* Herbicide Garlon\* 600 Herbicide Grazon\* DS Herbicide Lontrel\* 750SG Herbicide Starane\* 200 Herbicide Topik\* 240 Selective Herbicide Tordon\* 75-D Herbicide Recon® 520 Herbicide Tordon\*242 Cereal Herbicide Fusion\* WG Herbicide Select\* Herbicide Targa\* Herbicide Fusilade\* WG Herbicide Achieve\* Herbicide Verdict\* 520 Herbicide Precept\* Herbicide

# when mixing with atrazine water dispersible granule formulations, agitation is required.

In 2 Pro is compatible with cotton defoliant products such as: Thidiazuron, Ethephon.

In 2 Pro is compatible with the insecticides: Predator\* 300 Insecticide, Tracer\* Naturalyte Insect Control

Situations	Rate / 100 L of Water	Critical Comments
Cereals and Legumes	0.5 L – 1 L	When applying the herbicide Racer® or
	Rate recommendation will	Nitro®, apply at a rate of 1 L /100 L if
	vary depending upon the	targeting hard to control weeds.
	specific herbicide and the	When applying to a tankmix of Racer and
	prevailing conditions at the	Nitro, apply at a rate of 1 L /100 L.
	time of spraying.	When applying with the herbicide Precept* or
		Velocity* apply at a rate of 0.5 L /100 L.
Broadacre cropping including fallow (except	0.5 L	Applying a minimum of 250 mL of In2 Pro/
sugarcane)		per hectare.
Rice	1 L	Applying a minimum of 500 mL of In2 Pro/
Sugarcane	1L	per hectare.
Cotton Defoliation	0.5 L	Apply in combination with defoliants such as
		Dropp*, Mace® or Prep*
Spot and High Volume Spraying	0.5 L	Applying to the point of run-off.
Horticulture	0.5 L	Applying a minimum of 250 mL of In2 Pro/
		per hectare.

Important: Prior to use, read all directions on this label and on the herbicide (s) labels being used. The herbicide(s) labels for specific usage and restrictions have precedence over directions for use referred to on this label.

Note – 500 mL per 100 L is equivalent to 0.5% v/v, 1L per 100 L is equivalent to 1.0% v/v.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION