



Product Name: REBEL POUR ON FOR SHEEP
APVMA Approval No: 90703/129842

Label Name:	REBEL POUR ON FOR SHEEP
-------------	-------------------------

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	6 g/L ABAMECTIN
-------------------------	-----------------

Claims:	<p>For the control of sheep body lice (<i>Bovicola ovis</i>) on sheep off-shears, including lice resistant to synthetic pyrethroids.</p> <p>For the treatment and control of abamectin sensitive strains of gastrointestinal nematodes of sheep (including levamisole and benzimidazole resistant strains).</p> <p>When applied at the recommended dose rates to sheep off-shears (within 24 hours of shearing), REBEL POUR ON FOR SHEEP controls body lice (<i>Bovicola ovis</i>), including lice resistant to synthetic pyrethroids.</p> <p>REBEL POUR ON FOR SHEEP contains abamectin, a member of the macrocyclic lactone family of chemicals. It is effective against sensitive strains of the following internal parasites.</p> <p>REBEL POUR ON FOR SHEEP is for the treatment and control of the following abamectin sensitive adult and immature gastrointestinal nematodes (including levamisole and benzimidazole resistant strains):</p> <p>Barbers pole worm (<i>Haemonchus contortus</i>) including inhibited L4 and immature stages, Large stomach Worm (<i>Haemonchus placei</i>), Small brown Stomach Worm (<i>Teladorsagia circumcincta</i>), Stomach Hair Worm (<i>Trichostrongylus axei</i>), Black Scour Worm (<i>Trichostrongylus</i> spp), Nodule Worm (<i>Oesophagostomum columbianum</i>), Large Bowel Worm (<i>Oesophagostomum venulosum</i>).</p> <p>For best results REBEL POUR ON FOR SHEEP should be part of a total parasite control programme including internal and external parasites. Consult your veterinarian for a complete parasite control program.</p>
---------	---

RESISTANCE MAY DEVELOP TO ANY CHEMICAL

Net Contents:	2 L 2.5 L 5 L 6 L 10 L 15 L 20 L 22 L 25 L
---------------	--

Directions for Use:	
---------------------	--

Restrains:	DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption. DO NOT USE on unshorn sheep or shorn sheep more than 24 hours after shearing. DO NOT USE on lambs less than 10 kg bodyweight.
------------	---

Contraindications:	This product is contraindicated for use on lambs under 6 weeks of age or less than 10 kg body weight. This product should not be used on severely sick, debilitated or dying animals. This product should not be used on sheep that have not been cleanly shorn for any reason eg. Mycotic Dermatitis.
--------------------	--

Precautions:	Treat animals in accordance with good agricultural practice. Sheep should not be treated in extremely hot weather (> 40 °C) or if under heat stress.
--------------	--

Side Effects:	This product may irritate the skin of some sheep leading to visible treatment lines and a 'crusting' confined to the staple tip, which may still be evident at the following shearing. For further information please call ZAGRO AUSTRALIA CUSTOMER SERVICE on 1800 144 884.
---------------	--

Dosage and Administration:	<p>This product is for application to skin surface only, do not give orally or parenterally.</p> <p>REBEL POUR-ON FOR SHEEP is not recommended to be used off-shears on sheep shorn with 'cover' (snow) combs, as a reduction in efficacy is likely to occur.</p> <p>REBEL POUR-ON FOR SHEEP is not recommended to be used in ewes within 6 weeks of lambing or ewes with unshorn lambs at foot, as a reduction in lice efficacy is likely to occur.</p> <p>REBEL POUR-ON FOR SHEEP is not recommended to be used on housed (shedded) sheep that will not be exposed to natural sunlight for the first 2 months following treatment. The dye used in REBEL POUR-ON FOR SHEEP to identify treated sheep requires exposure to natural sunlight to degrade following treatment.</p> <p>The dose is 10 mL for each 10 kg of liveweight.</p> <p>Always dose on a liveweight basis. Dose the mob to the heaviest animal by liveweight in each group (ewes, rams, lambs), do not underdose. The heaviest animals in each group should be weighed. Check the accuracy of equipment before and during treatment.</p> <p>The following table may be used as a guide: Liveweight (kg) Dose (mL)</p>
----------------------------	--

	10 – 15 2 x 7.5 16 – 20 2 x 10 21 – 30 2 x 15 31 – 40 2 x 20 41 – 50 2 x 25 50 – 60 2 x 30 61 - 70* 2 x 35	Diagrams of applying product
	*add 10 mL for each kg above 70 kg.	
	Critical comments: Apply in two adjacent bands, either side of the backline, from the poll (bands may overlap along the neck) to the tail of each sheep using the recommended applicator.	

General Directions:	<p>NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p> <p>REBEL POUR ON FOR SHEEP is not recommended for use in species other than sheep. Use only in well-ventilated areas or outdoors. Close container when not in use. REBEL POUR ON FOR SHEEP is a short acting, non-residual product. Maximum lice kill at treatment and strict quarantine conditions for treated sheep are necessary for effective lice control. It is recommended that a thorough muster be conducted so that no untreated sheep remain. Boundary fences should be checked to ensure animals are contained and stray sheep are excluded. Treated sheep must be kept separate from lice infested sheep or sheep of unknown lice status. For effective lice control every sheep on the property should be treated at the same time.</p> <p>Efficacy of REBEL POUR ON FOR SHEEP against sheep body lice or abamectin sensitive nematodes is not adversely affected if applied when the skin is wet or if moderate rainfall occurs shortly after treatment. The antiparasitic activity of abamectin will be impaired if the formulation is applied to areas of skin with significant soiling with mud, dust, urine or faeces.</p> <p>The antiparasitic activity of abamectin will be impaired if the formulation is applied to roughly shorn sheep.</p> <p>2 Litre, 2.5 Litre, 5 Litre and 6 Litre Collapsible Packs:</p> <p>Use dosing equipment compatible with REBEL POUR ON FOR SHEEP. Follow manufacturer's directions for use and care of equipment. Other dosing equipment may be incompatible resulting in locking, incorrect dosage and leaking.</p>
---------------------	--

Withholding Periods:	<p>MEAT: DO NOT USE less than 21 days before slaughter for human consumption. MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption. WOOL HARVESTING INTERVAL: 6 weeks.</p>
----------------------	--

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 28 days before slaughter for export. Before using this product, confirm the current ESI from ZAGRO AUSTRALIA CUSTOMER SERVICE on 1800 144 884 or the APVMA website www.apvma.gov.au/residues</p> <p>Use of REBEL POUR-ON FOR SHEEP will not adversely affect trade in Australian wool providing the 6 week wool harvesting interval is observed.</p>
---------------	--

Safety Directions:	<p>Harmful if inhaled or swallowed. Do not inhale vapour. Will irritate eyes. Avoid contact with eyes. When using the product wear elbow-length chemical resistant gloves and goggles.</p>
--------------------	--

	If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.
First Aid Warnings:	
Additional User Safety:	<p>WOOL RE-HANDLING PERIOD: DO NOT re-handle treated sheep for 7 days unless wearing cotton overalls or equivalent clothing and chemical resistant gloves. Clothing must be laundered after each day's use.</p> <p>Additional information is listed in the Safety Data Sheet.</p>
Environmental Statements:	Very toxic to aquatic life. DO NOT contaminate streams, rivers, wetlands or watercourses with this product or used containers.
Disposal:	Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
Storage:	Store below 30 °C (room temperature) in the tightly closed original container in a cool dry place. Do not store for prolonged periods in direct sunlight.