



Product Name: Reflex-F Pour-On for Cattle
APVMA Approval No: 90929/137040

Label Name:	Reflex-F Pour-On for Cattle
-------------	-----------------------------

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS
------------------	--

Constituent Statements:	300 mg/mL TRICLABENDAZOLE 5 mg/mL ABAMECTIN Also contains: 500 mg/mL DIETHYLENE GLYCOL MONOBUTYL ETHER (solvent)
-------------------------	---

Claims:	<p>For the treatment and control of roundworms, liver fluke (all 3 stages) and external parasites of beef cattle.</p> <p>REFLEX-F POUR-ON FOR CATTLE contains abamectin, a member of the macrocyclic lactones family of chemicals. It is effective against sensitive strains of internal and external parasites. It also contains triclabendazole, a member of the benzimidazoles family of chemicals, and is effective against liver fluke. Resistance may develop to any chemical.</p> <p>INDICATIONS For the treatment and control of adult and immature stages of:</p> <p>Gastro-intestinal roundworms Barber's Pole Worm (<i>Haemonchus placei</i>) Small Brown Stomach Worm (<i>Ostertagia ostertagi</i>) (including inhibited stages) Small Intestinal Worm (<i>Cooperia oncophora</i>, <i>Cooperia punctata</i>) Stomach Hair Worm (<i>Trichostrongylus axei</i>) Hookworm (<i>Bunostomum phlebotomum</i>) Large Bowel Worm (<i>Oesophagostomum radiatum</i>) Whipworm (<i>Trichuris</i> spp.) Thin Necked Intestinal Worm (<i>Nematodirus spathiger</i>) Intestinal Threadworm (<i>Strongyloides papillosus</i>)</p> <p>Lungworm (<i>Dictyocaulus viviparus</i>)</p> <p>Liver fluke (<i>Fasciola hepatica</i>) including adult (12 week old), immature (6 week old) and early immature (4 week old) larvae</p> <p>Sucking and biting lice Long Nosed Sucking Lice (<i>Linognathus vituli</i>) Short Nosed Sucking Lice (<i>Haematopinus eurysternus</i>)</p>
---------	--

	<p>Biting Lice (<i>Bovicola bovis</i>)</p> <p>Little Sucking Lice (<i>Solenopotes capillatus</i>)</p> <p>Also aids in the control of:</p> <p>Buffalo Fly (<i>Haematobia irritans exigua</i>)</p> <p>Cattle Tick (<i>Rhipicephalus microplus</i>) (Not recommended in strategic dipping programs for cattle tick control)</p>
--	--

Net Contents:	1 L, 2.5 L, 3 L, 5 L, 5.5 L, 10 L, 15 L
---------------	---

Directions for Use:	
---------------------	--

Restrains:	DO NOT USE on cows which are producing or may in the future produce milk that may be used or processed for human consumption.
------------	---

Contraindications:	This product is contraindicated for use in calves less than 50 kg bodyweight.
--------------------	---

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	<p>The dose rate is 1 mL/10 kg body weight.</p> <p>Apply along the backline of the animal, using a standard pour-on applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected.</p> <p>Dosage chart Liveweight (kg); Dose volume (mL)</p> <p>51-75; 7.5 76-100; 10 101-125; 12.5 126-150; 15 151-175; 17.5 176-200; 20 201-250; 25 251-300; 30 301-350; 35 351-400; 40 401-450; 45 451-500; 50 501-550; 55 551-600; 60 601-650; 65</p> <p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION</p> <p>Cattle heavier than 650 kg should be dosed at 1 mL per 10 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not</p>
----------------------------	--

	underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing.
General Directions:	<p>REFLEX-F POUR-ON is rainfast. Rain occurring as early as 2 hours after treatment will not affect the efficacy of the product.</p> <p>A white deposit may form on the back of treated animals. This is quite normal. It is not a sign of treatment failure and should not be interfered with. Do not brush off or wash off. This deposit may last for a few days or weeks depending on weather conditions.</p> <p>Temporary skin reaction may occur along the application site, especially if applied during very hot weather. This should resolve without treatment. If signs persist or worsen, contact your veterinarian.</p> <p>After use clean all equipment with cooking oil. Do not use water.</p>
Withholding Periods:	<p>MEAT: DO NOT USE less than 49 days before slaughter for human consumption. Beef calves born to cows that were treated with REFLEX-F POUR-ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for human consumption for 49 days after treatment of the dam.</p> <p>MILK: DO NOT USE in cows which are producing or may in the future produce milk that may be used or processed for human consumption.</p>
Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 140 days before slaughter for export. Before using this product, confirm the current ESI from Alleva Animal Health on 1-800 ALLEVA (1-800 255 382) or the APVMA website (www.apvma.gov.au/residues).</p> <p>Beef calves born to cows that were treated with REFLEX-F POUR-ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for the export market for 140 days after treatment of the dam.</p>
Safety Directions:	Harmful if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash goggles and contaminated clothing.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.
First Aid Warnings:	
Additional User Safety:	Additional information is listed in the safety data sheet.
Environmental Statements:	<p>ENVIRONMENTAL PROTECTION</p> <p>Very toxic to aquatic life. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.</p>
Disposal:	[1 L] Dispose of container by wrapping with paper and putting in garbage.

[Other pack sizes] Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals onsite. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Storage:

Store below 30°C (Room Temperature).