



Product Name: BALISTIK POUR-ON FOR CATTLE
APVMA Approval No: 92358/140931

Label Name:	BALISTIK POUR-ON FOR CATTLE
-------------	-----------------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	15 g/L FLUAZURON 5 g/L IVERMECTIN
-------------------------	--------------------------------------

Claims:	<p>For strategic control of fluazuron susceptible cattle tick, <i>Rhipicephalus microplus</i>, including strains resistant to organophosphates, synthetic pyrethroids and amidines, on beef cattle.</p> <p>For the treatment and control of ivermectin sensitive internal and external parasites of beef cattle.</p> <p>BALISTIK POUR-ON FOR CATTLE contains fluazuron and ivermectin, a member of the avermectin (macrocyclic lactone) family. It is effective against cattle ticks and ivermectin sensitive internal and external parasites of beef cattle.</p> <p>Resistance may develop to any chemical.</p> <p>Cattle tick: <i>Rhipicephalus microplus</i></p> <p>Gastrointestinal roundworms:</p> <p>For the control of adult and immature (L4): Small brown stomach worms (including inhibited stages) <i>Ostertagia ostertagi</i> Barber's pole worm <i>Haemonchus placei</i> Stomach hair worm <i>Trichostrongylus axei</i> Intestinal hair worm <i>Trichostrongylus colubriformis</i> Small intestinal worm <i>Cooperia</i> spp. Nodule worm <i>Oesophagostomum radiatum</i> <i>Oesophagostomum venulosum</i></p> <p>And adult stages of: Whipworm</p>
---------	--

	<p>Trichuris spp. Intestinal threadworm Strongyloides papillosus</p> <p>And immature stages of: Thin necked intestinal worm L4 Nematodirus spp. Lungworm (adult and immature) Dictyocaulus viviparus Eyeworm (adult stages) Thelazia spp.</p> <p>Sucking and biting lice: Linognathus vituli Haematopinus eurysternus Bovicola (Damalinia) bovis Solenopotes capillatus</p> <p>Mites: Chorioptes bovis Sarcoptes scabiei var bovis</p> <p>Buffalo flies: Haematobia irritans exigua</p> <p>Persistent activity: BALISTIK POUR-ON FOR CATTLE applied at the recommended dose of 5 mL/50 kg liveweight controls infestations of Haematobia irritans exigua (buffalo fly) for up to 7 days after treatment.</p>
--	--

Net Contents:	1 L, 2.5 L, 5 L, 10 L, 20 L, 25 L
---------------	-----------------------------------

Directions for Use:	
---------------------	--

Restrictions:	DO NOT USE on cows which are producing or may in the future produce milk that may be used or processed for human consumption.
---------------	---

Contraindications:	This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	<p><Table > A representative sample of animals should be weighed before treatment either with scales or a weigh band. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not underdose. Where there is a large variation in size within the group, draft into 2 or more lines based on body weight, to avoid excessive overdosing.</p> <p>Critical comments: 1. Retreat when you first see ticks but not before at least 6 weeks after the previous treatment. 2. Maximum number of treatments per annual tick season: 3.</p>
----------------------------	---

3. To achieve optimal efficacy, BALISTIK POUR-ON FOR CATTLE relies on a combination of absorption through the skin and oral ingestion (via grooming). For oral ingestion it is critical that the product is applied as 2 x 7 cm wide bands, from the shoulder to the rump, that are well separated on either side of the spine. Use a manufacturer approved applicator. DO NOT use any other applicator.
4. The weight of cattle should be determined by either scales or a weighband. Dose individuals according to the heaviest animal in the mob. Recommended rate gives 1.5 mg fluazuron and 0.5 mg ivermectin per kg bodyweight of beef cattle. DO NOT underdose.
5. There is no need to treat calves for cattle tick as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves.
6. BALISTIK POUR-ON FOR CATTLE should not be applied to show cattle as a light scurfiness may be visible for some time after treatment.
7. The effect of BALISTIK POUR-ON FOR CATTLE is not adversely affected if rain occurs after application.

This section contains file attachment.

General Directions:	<p>Cleaning Instructions After use wash BALISTIK POUR-ON FOR CATTLE applicator with detergent and hot water. Dispose of the rinsates in a disposal pit away from desirable plants and their roots and watercourses.</p> <p>BALISTIK POUR-ON FOR CATTLE Treated Areas - Introduction of New Cattle Tick infested cattle introduced onto properties where a BALISTIK POUR-ON FOR CATTLE treatment regime is in place should be treated with BALISTIK POUR-ON FOR CATTLE prior to release.</p> <p>Handling of Treated Animals Wear protective clothing recommended under Safety Directions, if handling treated animals within 24 hours of application.</p> <p>To avoid operator contact with the BALISTIK POUR-ON FOR CATTLE when concurrent treatments are carried out, BALISTIK POUR-ON FOR CATTLE should be applied last. For local tick control recommendations contact your local Department of Agriculture.</p> <p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p>
---------------------	--

Withholding Periods:	MEAT: DO NOT USE less than 28 days before slaughter for human consumption. Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.
----------------------	---

Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from Alleva Animal Health on 1-800 ALLEVA (1-800 255 382) or the APVMA website (apvma.gov.au/residues).
---------------	--

Safety Directions:	Harmful if swallowed. May irritate the skin. Will irritate the eyes. Avoid contact with eyes and skin. When preparing product for use and using product, wear elbow-length PVC gloves. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves.
--------------------	---

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	Additional information is listed in the safety data sheet.
-------------------------	--

Environmental Statements:	ENVIRONMENTAL PROTECTION The product is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.
---------------------------	--

Disposal:	<p>[1 L]: Dispose of container by wrapping with paper and putting in garbage.</p> <p>[2.5 L, 5 L, 10 L, 20 L and 25 L]: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals onsite. If recycling, replace cap and return clean container to recycler or designated collection point. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately at the drumMUSTER collection site. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
-----------	---

Storage:	Store below 30°C (room temperature). Store in original container tightly closed in a dry, cool place. Store out of direct sunlight.
----------	---

Dosage and Administration

Animals and Pests	Bodyweight (kg)	Dose (mL)
Cattle Tick (<i>Rhipicephalus microplus</i>) Ivermectin-sensitive gastrointestinal roundworms (including inhibited <i>Ostertagia ostertagi</i>) Lungworms Sucking and biting lice Sarcoptic and chorioptic mites Buffalo fly (<i>Haematobia irritans exigua</i>)	101 – 150	15
	151 – 200	20
	201 – 250	25
	251 – 300	30
	301 – 350	35
	351 – 400	40
	401 – 450	45
	451 – 500	50
	501 – 550	55
	551 – 600	60
601 – 650	65	
	Over 650	5 mL/50 kg