



Product Name: Covine Snap Extra Spray On Sheep Blowfly Treatment  
APVMA Approval No: 93561/139541

Label Name:	Covine Snap Extra Spray On Sheep Blowfly Treatment
-------------	--

Signal Headings:	FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	65 g/L DICYCLANIL
-------------------------	-------------------

Claims:	<p>For the protection of sheep against fly strike (caused by dicyclanil-susceptible strains of blowflies (<i>Lucilia cuprina</i>)). When applied as directed this product will protect against fly strike for up to 29 weeks. For the protection of mulesing and marking wounds on sheep against fly strike (caused by dicyclanil-susceptible strains of blowflies (<i>Lucilia cuprina</i>)) during the wound healing process.</p> <p>SNAP EXTRA Spray On contains dicyclanil, a member of the pyrimidine group of chemicals. It is effective against strains of <i>Lucilia cuprina</i> that are sensitive to dicyclanil. It is not as effective against strains of <i>Lucilia cuprina</i> that develop resistance to dicyclanil.</p> <p>Resistance may develop to any chemical. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance.</p>
---------	---

Net Contents:	800 mL, 5 L, 20 L, 25 L
---------------	-------------------------

Directions for Use:	
---------------------	--

Restraints:	<p>DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.</p> <p>DO NOT USE on sheep less than 4 weeks after shearing.</p> <p>DO NOT apply at the same time as any other backline product.</p> <p>Re-treatment interval: DO NOT re-treat animals for a minimum of 9 weeks after last treatment.</p>
-------------	--

Contraindications:	
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	This section contains file attachment.
----------------------------	--

General Directions:	<p><b>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION</b></p> <p>SNAP EXTRA Spray-On provides season long protection to sheep against body, breech and poll strike for up to 29 weeks and may be used on sheep with any length wool from 4 weeks after shearing.</p> <p>SNAP EXTRA Spray-On also protects mulesing and marking wounds on lambs against fly strike during the wound healing process. The dose volume is determined by the weight of sheep being treated. Apply SNAP EXTRA Spray-On according to the diagrams and tables in the Directions for Use. Apply directly from the container or an appropriate back pack using a spray-on applicator gun recommended by the manufacturer. SNAP EXTRA Spray-On may be applied to all breeds of sheep to obtain protection against fly strike. SNAP EXTRA Spray-On is applied as a low-volume treatment onto areas where protection is required.</p> <p>When applying SNAP EXTRA Spray-On, the nozzle of the applicator gun should be high enough above the fleece (approximately 15-25 cm), to achieve a 15 cm wide band at each pass for protection from body strike. For treatment of mulesing wounds, the band must be wide enough to cover the wound and 25 mm onto the surrounding wool.</p> <p>The product should not be applied during gusty or windy weather as spray drift may reduce the period of protection. It is good agricultural practice for mulesing/marking to be carried out at times of minimal fly activity. Reduced efficacy may occur in sheep affected with dermatophilosis. Significant soiling with dags, mud, dust or urine may reduce efficacy.</p> <p>Where significant dags are present, a light crutching is recommended prior to treatment. Protection period may be reduced if sheep scour heavily following application.</p> <p>Not to be mixed with any other product.</p> <p>Not to be diluted.</p>
---------------------	---

Withholding Periods:	<p>MEAT: DO NOT USE less than 14 days before slaughter for human consumption.</p> <p>WOOL: DO NOT USE less than 3 months before shearing or fibre collection.</p>
----------------------	---

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 9 weeks before slaughter for export. Before using this product, confirm the current ESI from Nutrien Ag Solutions Limited on 1800 448 892 or the APVMA website (<a href="http://www.apvma.gov.au/residues">www.apvma.gov.au/residues</a>).</p> <p>Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable</p>
---------------	---

	residues; adequate treatment records should be kept and made available, if requested by wool buyers.
Safety Directions:	May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product wear elbow length rubber gloves. Wash hands after use.
First Aid Instructions:	First aid is not generally required. If in doubt, contact a doctor or the Poisons Information Centre: Phone Australia 131 126.
First Aid Warnings:	
Additional User Safety:	Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling. If additional hazard information is required, refer to the Safety Data Sheet.
Environmental Statements:	<b>ENVIRONMENTAL PROTECTION</b> Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.
Disposal:	Containers <1L: Dispose of container by wrapping with paper and putting in garbage.  Containers >1L: Triple-rinse container. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.  If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately at the drumMUSTER collection site.
Storage:	<b>KEEP OUT OF REACH OF CHILDREN.</b> Store below 30 °C (room temperature). Do not freeze, protect from light. Store in original container tightly closed in a dry, cool place.

**DOSAGE AND ADMINISTRATION:****SHAKE WELL BEFORE USE**

Use remaining product within 12 months of opening the container.

**FOR THE PREVENTION OF BODY STRIKE (Diagram 1)**

SHEEP WEIGHT (KG)	TOTAL VOLUME/SHEEP (ML)	VOLUME PER TREATMENT BAND (ML)	NO. OF BANDS
10-20	10	10	1
21-30	12	12	1
31-50	15	7.5	2
>50	18	9	2

**FOR THE PREVENTION OF BREECH STRIKE (Diagram 1)**

SHEEP WEIGHT (KG)	TOTAL VOLUME/SHEEP (ML)	VOLUME PER TREATMENT BAND (ML)	NO. OF BANDS
10-20	10	10	1
21-30	12	12	1
31-50	15	7.5	2
>50	18	9	2

**FOR THE PREVENTION OF POLL STRIKE (Diagram 1)**

- Rams and sheep susceptible to poll strike should be treated. Apply 5mL of SNAP EXTRA Spray-On to the areas requiring protection, around the poll and horns.

**FOR THE PREVENTION OF FLY STRIKE AFTER MULESING (Diagram 2)**

Apply two bands, one over each side of the mulesing wound and overlapping 25 mm onto the surrounding wool.

SHEEP WEIGHT (KG)	TOTAL VOLUME/SHEEP (ML)	VOLUME PER TREATMENT BAND (ML)	NO. OF BANDS
Up to 30	8	4	2
31-50	10	5	2

**FOR THE PREVENTION OF FLY STRIKE AFTER MARKING (Diagram 3)**

- When a tail has been amputated, apply 4 mL over the crutch ensuring sufficient coverage of the open wound.
- When the tail has not been amputated, ensure sufficient coverage of the tail where the rubber ring has been applied.


In the case of ram lambs a further 4 mL can be sprayed over and around the scrotum area after the rubber ring has been applied.

SHEEP WEIGHT (< 30 KG)	TOTAL VOLUME/SHEEP (ML)	VOLUME PER TREATMENT BAND (ML)	NO. OF BANDS
Ram lambs	8	4	2
Ewe lambs	4	4	1

**CRITICAL COMMENTS**

1. Use only a spray-on application gun.
2. Ensure the applicator gun is clean.
3. Calibrate applicator gun using an accurate volume measure.
4. Ensure all areas to be protected are treated.
5. When preventing body strike ensure bands overlap along the mid-line (as in diagram 1).
6. When protecting from mulesing wounds from fly strike, take special care to treat the wound and the surrounding 25 mm wool (as in diagram 2).
7. When protecting marking wounds from fly strike, take special care to treat the wound and surrounding wool (as in diagram 3).
8. This product is not recommended for treating existing strikes. Use a product registered as a dressing for existing strikes.
9. Heavy rain following the application of this product could diminish the period of protection.
10. SNAP EXTRA Spray-On contains a scourable colouring agent for easy band identification.
11. Withholding periods must be observed following application of SNAP EXTRA Spray-On for any purpose including after mulesing and/or marking as finite residues will occur.

**Application Diagrams**



**Diagram 1**



**Diagram 2**



**Diagram 3**